

Références Bibliographiques

- Ackley S.F., Sullivan C.W., 1994. Physical controls on the development and characteristics of Antarctic sea ice biological communities - a review and synthesis. *Deep-Sea Res.*, 41, 1583-1604.
- Aksnes D.L., Wassmann P., 1993. Modeling the significance of zooplankton grazing for export production. *Limnol. Oceanogr.*, 35, 978-985.
- Alcaraz M., 1988. Summer zooplankton metabolism and its relation to primary production in the Western Mediterranean. *Oceanol. Acta*, 9 (Spec. Issue), 185-191.
- Alcaraz M., Saiz E., Fernandez J.A., Trepot I., Figueiras F., Cablet A., Bautista B., 1998. Antarctic zooplankton metabolism: carbon requirements and ammonium excretion of salps and crustacean zooplankton in the vicinity of the Bransfield Strait during January 1994. *J. Mar. Syst.*, 17, 347-359.
- All dredge A.L., Gotschalk G.C., 1989. Direct observation of the mass flocculation of diatom blooms: characteristics, settling velocities and formation of diatom aggregates. *Deep-Sea Res.*, 36, 159-172.
- Allain C., 1960. Topographie dynamique et courants généraux dans le bassin occidental de la Méditerranée. *Rev. Trav. Inst. Pêches Marit.*, 4, 27-42.
- Allen K.R., 1966. Some method for estimating exploited populations. *J. Fish. Res. Bd. Canada*, 32, 1123-1143.
- Andersen V., Nival P., 1988. Modèles d'écosystème pélagique des eaux côtières de la Mer Ligure. *Oceanol. Acta*, 9 (Spec. Issue), 211-217.
- Andersen V., Nival P., 1991. A model of the diel vertical migration of zooplankton based on euphausiids. *J. Mar. Res.*, 49, 153-175.
- Andersen V., Nival P., Harris R.P., 1987. Modelling of a planktonic ecosystem in an enclosed water column. *J. mar. biol. Ass. UK*, 67, 407-430.
- Andersen V., Rassoulzadegan F., 1991. Modèle vertical de l'écosystème pélagique marin. Réseau microbien et sédimentation des particules biogéniques. *J. Rech. Océanogr.*, 16, 16-22.
- Andreas E.L., Ackley S.F., 1982. On the differences in ablation seasons of Arctic and Antarctic sea ice. *J. Atmos. Sci.*, 39, 440-447.
- Andreas E.L., Tucker W.B., Ackley S.F., 1984. Atmospheric boundary-layer modification, drag coefficient, and surface heat flux in the Antarctic marginal ice zone. *J. Geophys. Res.*, 89, 649-661.
- Antia N.J., McAllister C.D., Parsons T.R., Stephens K., Strickland J.D.H., 1963. Further measurements of primary production using a large-volume plastic sphere. *Limnol. Oceanogr.*, 8, 166-183.
- Arrigo K.R., McClain C.R., 1994. Spring phytoplankton production in the Western Ross Sea. *Science*, 266, 261-263.
- Arrigo K.R., Robinson D.H., Worthern D.L., Dunbar R.B. DiTullio G.R., VanWoert M., Lizotte M.P., 1999. Phytoplankton community structure and the drawdown of nutrients and CO₂ in the Southern Ocean. *Science*, 283, 365-367.
- Arrigo K.R., Weiss A.M., Smith W.O.Jr., 1998a. Physical forcing of phytoplankton dynamics in the southwestern Ross Sea. *J. Geophys. Res.*, 103(C1), 1007-1021.
- Arrigo K.R., Worthern D.L., Lizotte M.P., Dixon P., Dieckmann G., 1997. Primary Production in Antarctic Sea Ice. *Science*, 276, 394-397.
- Arrigo K.R., Worthern D., Schnell A., Lizotte M.P., 1998b. Primary production in Southern Ocean waters. *J. Geophys. Res.*, 103, 15587-15600.
- Atkinson A., Ward P., Williams R., Poulet S.A., 1992. Feeding rates and diel vertical migration of copepods near South Georgia: comparison of shelf and oceanic sites. *Mar. Biol.*, 114, 49-56.
- Azam F., 1998. Microbial control of oceanic carbon flux: the plot thickens. *Science*, 280, 694-696.

- Azam F., Ammermann J.W., 1984. Cycling of organic matter by bacterioplankton in pelagic marine ecosystems: microenvironmental considerations. In M.J.R. Fasham (Ed): Flow of energy and materials in marine ecosystems, Plenum, pp 345-360.
- Azam F., Fenchel T., Field J.G., Gray J.S., Meyer-Reil L.A., Thingstad F., 1983. The ecological role of water-column microbes in the sea. Mar. Ecol. Prog. Ser., 10, 257-263.
- Azam F., Smith D.C., 1991. Bacterial influence on the variability in the ocean's biogeographical state: a mechanistic view. In S. Demers (Ed): Particle analysis in oceanography, Springer Verlag, Berlin, pp 213-236.
- Azzali M., Kalinowski J., 1999. Spatial and temporal distribution of krill *Euphausia superba* biomass in the Ross Sea (1989-1990 and 1994). In F. Faranda, L. Guglielmo and A. Ianora (Eds): Ross Sea Ecology. Springer, Berlin Heidelberg, pp. 433-455.
- Bainbridge, R., 1961. Migrations. In Waterman (Ed): The physiology of Crustacea. Academic Press, Lond. New York, 431-463.
- Banse K., 1977. Determining the carbon-to-chlorophyll ratio of natural phytoplankton. Mar. Biol., 41, 199-212.
- Bavestrello, G., Cattaneo-Vietti R., Sara M., Veeschekens C., Hecq J.H., 1992. Chromatographic analysis on sponges with autotrophic symbionts. Rapp. Comm. int. Mer Médit., 33, 30.
- Bay D., 1978. Etude in situ de la production primaire d'un herbier de posidonies (*Posidonia oceanica* (L) Delile) de la baie de Calvi, Corse. Ph.D. Thesis, Univ. Liège, Belgium, 251 pp.
- Beckers, O., Nihoul, J.C.J. R. Wollast, 1976. La circulation résiduelle et la caractérisation des masses d'eau dans la zone belge. In J.C.J. Nihoul (Editeur.). Modélisation des systèmes marins, Projet Mer, Rapport final, Services du Premier Ministre, Programmation de la Politique scientifique, Bruxelles, 95-130.
- Beckers J.M., 1991. Application of the GHER 3D general circulation model to the Western Mediterranean. J. Mar. Syst., 1, 315-332.
- Beers J.R., 1966. Studies on the chemical composition of the major zooplankton groups in the Sargasso Sea of Bermuda. Limol. Oceanogr., 11, 520-528.
- Beklemishev C.W., 1962. Superfluous feeding of marine herbivorous zooplankton. Rapp. P.-V. Reun. Cons. Perm. Int. Explor. Mer. 153, 108-113.
- Belkhiria S., Goffart A., Hecq J.H., 1996. Determination of zooplankton grazing pressure and faecal pellet production in a frontal zone of the North Sea. Progress in Belgian Oceanographic Research, 19-22.
- Berland B.R., Benzhitski A.G., Burla-Kova Z.P., Georgieva L.V., Izmestieva M.A., Kholodov V.I., Maestrini S.Y., 1988. Conditions hydrologiques estivales en Méditerranée, répartition du phytoplancton et de la matière organique. Oceanol. Acta, 9 (Spec. Issue), 163-177.
- Berner A., 1962. Feeding and respiration in the copepod *temora longicornis* (Mueller) J. mar. biol. Ass. U.K., 42, 625-640.
- Béthoux J.P., 1979. Budgets of the Mediterranean Sea. Their dependence on the local climate and on the characteristics of the Atlantic waters. Oceanol. Acta, 2, 157-163.
- Béthoux J.P., 1981. Le phosphore et l'azote en mer Méditerranée, bilans et fertilité potentielle. Mar. Chem., 10, 141-158.
- Béthoux J.P., Prieur L., 1979. Evaluation des flux d'eau de la circulation du Nord-Est du bassin occidental. Rapp. Comm. int. Mer Médit.. 25-26, 67-68.
- Béthoux J.P., Prieur L., 1983. Hydrologie et circulation en Méditerranée nord-occidentale. Pétrole et technique, 299, 25-34.
- Béthoux J.P., Prieur L., Nyffeler F., 1981. Utilisation de moyennes hydrologiques pour le calcul des flux d'eaux dans le bassin Liguro-Provençal. Rapp. Comm. int. Mer Médit., 27, 6, 193-196.
- Bidigare R.R., Frank T. J., Zastrow C., Brooks J.M., 1986. The distribution of algal chlorophylls and their degradation products in the Southern Ocean. Deep-Sea Res., 33, 923-937.
- Bidigare R.R., Kennicutt M.C., Brooks J.M., 1985. Rapid determination of chlorophylls and their degradation products by high-performance liquid chromatography. Limnol. Oceanogr., 30, 432-435.

- Bidigare R.R., Marra J., Dickey T.D., Iturriaga R., Baker K.S., Smith R.C., Pak H., 1990. Evidence for phytoplankton succession and chromatic adaptation in the Sargasso Sea during spring 1985. *Mar. Ecol. Prog. Ser.*, 60, 113-122.
- Billen G., Bossicart M., Daro M.H., Deboever R., Deconinck L., Govaere J., Hecq J.H., Heip C., Hovart P., Janssens D., Joiris C., Mommaerts J.-P., Pichot, G., Redant F., Thielemans L., Runfola Y., Van Beveren C., VanDamme O., Van DeVelde, J., 1976. Chaînes trophiques et cycle des nutriments. In J.C.J. Nihoul et F. Polk (Eds): Rapport final, Programme National "Recherche et Développement, Environnement - Projet mer de la Pol. Scientifique, 8, 340 pp.
- Billen G., Lancelot C., De Becker E., Servais P., 1988. Modelling microbial processes (phyto- and bacterioplankton) in the Schelde estuary. *Hydrobiological Bulletin*, 22, 43-55.
- Billen G., Joiris C., Meyer-Reil L., Lindeboom H., 1990. Role of bacteria in the North Sea ecosystem. *Neth. J. of Sea Res*, 26 (2-4), 265-293.
- Blackman F.F., 1905. Optima and limiting factors. *Ann. Bot.*, 19, 281-295.
- Blackbourn D.J., Taylor F.R.J., Blackbourn J., 1973. Foreign organelle retention by ciliate. *J. Protozool.*, 20, 286-288.
- Bolin B. 1983. Changing global geochemistry. In P. Brewer (Ed): *Oceanography. The present and future*. Springer - Verlag, 305-326.
- Boucher J., 1988. Space-time aspects in the dynamics of plankton stages. In B.J. Rothschild (Editor), *Towards a theory on biological-physical interactions in the world ocean*. Kluwer Academic Publishers, the Netherlands, 203-214.
- Boucher J., Ibanez F., Prieur L., 1987. Daily and seasonal variations in the spatial distribution of zooplankton populations in relation to the physical structure in the Ligurian Sea Front. *J. Mar. Res.*, 45, 133-173.
- Bougis P., 1974. *Ecologie du plancton marin, II. Le zooplancton*. Masson, 196 pp.
- Bouquegneau J.M., Billen G., Joiris C., Daro M.H., Lancelot C., Hecq J.H., Bertels A., Bossicart M., Somville M., Mommaerts J.P., Wijnant J., Wehren J.C., 1982. Carbon cycling in the Belgian coastal zone and adjacent areas. I. General structure of the ecosystem. *Int. Counc. Explor. Sea. Biol. Oceanogr. Comm.*, C.M. 1982/L, 31, 5-35.
- Bouquegneau J.M., Gieskes W.W.C., Kray G.W., Larsson A.M., 1992. Influence of physical and biological processes on the concentration of O₂ and CO₂ in the ice-covered Weddell Sea in the spring of 1988. *Polar Biol.*, 12, 163-170.
- Braconnor J.C., 1967. Sur la possibilité d'un cycle court de développement chez le tunicier pélagique *Doliolum nationalis* Borg. *C. R. Acad Sci. Paris*, 264, 1434-1437.
- Bratbak G., Egge J.K., Heldal M., 1993. Viral mortality of the marine alga *Emiliania huxleyi* (Haptophyceae) and the termination of algal blooms. *Mar. Ecol. Prog. Ser.*, 93, 39-48.
- Brockmann U.H., Laane R.W., Postma H., 1990. Cycling of nutrient elements in the North Sea. *Neth. J. of Sea Res*, 26 (2-4), 239-264.
- Broecker W.S., 1987. The biggest chill. *Nat Hist. Mag.*, 97, 74-82.
- Brooker R.E., 1971. Fine structure of *Bodo saltans* and *Bodo caudatus* *Zoomastigophora, Protozoa*. *Bull. Brit. Mus. Lond. Zoology*, 22, 87-102.
- Brooks J.E., 1964. Acclimation to light intensity in two species of marine phytoplankton *Dunaliella tertiolecta* Butcher and *Skeletonema costatum* (Grev.) Cleve. *M.S. Thesis University of Southern California, Los Angeles* 182pp.
- Brohée M., Goffart A., Frankignoulle M., Henri V., Mouchet A., Hecq J.H., 1989. Variations printanières des communautés planctoniques en baie de Calvi (Corse) en relation avec les contraintes physiques locales. *Cah. Biol. Mar.*, 30, 321-328.
- Cadée G.C., 1985. Macroaggregates of *Emiliania huxleyi* in sediment trap. *Mar. Ecol. Prog. Ser.*, 24, 193-196.
- Cadée G.C., 1986. Organic carbon in the water column and its sedimentation, Fladden Ground (North Sea), May 1983. *Neth. J. of Sea Res*, 20, 347-358.
- Cadée G.C., 1992. Trends in Marsdiep phytoplankton. In N. Danker, C.J. Smit, M. Scholl (Eds): *Present and future conservation of the Wadden Sea. Proceedings of the International Wadden Sea Symposium. Armeland*. 1990. *Neth. Inst. Sea Res. Publ. Ser.*, 20, 143-149

- Cahet G., Fiala M., Jacques G., Panouse M., 1972. Production primaire au niveau de la thermocline en zone néritique de la Méditerranée Nord-Occidentale. Mar. Biol., 14, 32-40.
- Campbell L., Nolla H.A., Vaulot D., 1994. The importance of Prochlorococcus to community structure in the central North Pacific Ocean. Limnol. Oceanogr., 39, 954-961.
- Carlotti F., Giske J., Werner F., 2000. Modeling zooplankton dynamics. In ICES Zooplankton Methodology Manual. Academic Press, 571-668.
- Carrada G.C., Hopkins T.S., Bonaduce G., Ianora A., Marino D., Modigh M., Ribera D'Alcalà M., Scotti di Carlo B., 1980. Variability in the hydrographic and biological features of the Gulf of Naples. Mar. Ecol., 1, 105-120.
- Carrada G.C., Fresi E., Marino D., Modigh M., Ribera D'Alcalà M., 1981. Structural analysis of winter phytoplankton in the Gulf of Naples. J. Plankton Res., 3, 291-314.
- Carrada G.C., Mangoni, O., Russo G.F., Saggiomo V., 1999. Phytoplankton size-fractionated biomass in the Ross sea: Spatial and temporal variations during the austral spring. In F. Faranda, L. Guglielmo & A. Ianora (Eds): Ross Sea Ecology. Springer, Berlin Heidelberg, pp. 205-216.
- Catalano G., Benedetti F., Goffart A., Iorio M., 1991. Distribution of dissolved oxygen, ph, total alkalinity and nutrients in the Southern Ocean and the Ross Sea (R/V "Cariboo" 1989-1990 cruise). In: National Scientific Commission for Antarctica, Oceanographic Campaign 1989-1990. Data Report Part I, pp. 11-23.
- Catalano G., Povero P., Fabiano M., Benedetti F., Goffart A., 1997. Nutrient utilisation and particulate organic matter changes during summer in the upper mixed layer (Ross Sea, Antarctica). Deep-Sea Res., 44, 97-112.
- Champalbert G., 1969. Microdistribution d'un Pontellidae dans le Golfe de Marseille: *Anomalocera patersoni*. Mar. Biol., 2, 346-349.
- Chapelle S., Gilles R., Hecq J.H., 1980. Phospholipids, fatty acids, free aminoacids and proteic aminoacids in zooplankton from the straight of Dover (North Sea). In R. Gilles (Ed), Europ. Soc. compar. Physiol. Biochem, 177-178.
- Checkley D.M.Jr., Uye S., Dagg M.J., Mullin M.M., Omori M., Onbe T., Zhu M., 1992. Diel variation of the zooplankton and its environment at neritic stations in the Inland Sea of Japan and the north-west Gulf of Mexico. J. Plankton Res., 14, :1-40.
- Chisholm S.W., Olson R.J., Zettler E.R., Goericke R., Waterbury J.B., Welschmeyer H.A., 1988. A novel free-living prochlorophyte abundant in the oceanic euphotic zone. Nature, 334, 340-343.
- Chisholm S.W., Morel F.M.M., 1991. What controls phytoplankton production in nutrient-rich areas of the open sea? Limnol. Oceanogr. 36, 1507-1965.
- Cho B.C., Azam F., 1990. Biogeochemical significance of bacterial biomass in the ocean's euphotic zone. Mar. Ecol. Prog. Ser., 63, 253-259.
- Clarke A.J., 1985. Energy flow in the Southern Ocean Food Web. In W.R. Sigfried, P.R. Condy & R.M. Laws (Eds): Antarctic Nutrient Cycles and Food Webs, Springer-Verlag, Berlin Heidelberg New York, pp. 573-580.
- Clarke A.J., Leakey R.J.G., 1996. The seasonal cycle of phytoplankton, macronutrients, and the microbial community in a nearshore Antarctic marine ecosystem. Limnol. Oceanogr., 41, 1281-1294.
- Claustre H., Poulet S.A., Williams R., Marty J.C., Coombs S., Ben Mijh F., Hapette A.M., Martin-Jezequel V., 1990. A biochemical investigation of a *Phaeocystis* sp. Bloom in the Irish Sea. J. Mar. Biol. Assoc. U.K., 70, 197-207.
- Claustre H., Marty J.C., Kerhervé P., Prieur L., Hecq J.H., 1992. Distributions verticales et horizontales des marqueurs pigmentaires phytoplanctoniques durant la campagne Almofront-1. In A. Sournia & L. Prieur (Eds): Processes and budgets in geostrophic fronts. (France JGOFS), 35-36.
- Claustre H., Kerhervé P., Marty J.C., Prieur L., Videau C., Hecq J.H., 1994. Phytoplankton dynamics associated with a geostrophic front: Ecological and biogeochemical implications. J. Mar. Res., 52, 711-742.

- Cloern J.E., Grenz C., Vidergar-Lucas L., 1995. An empirical model of the phytoplankton chlorophyll: carbon ratio - the conversion between productivity and growth. Limnol. Oceanogr., 7, 1310-1313.
- Clutter R.I., 1969. The microdistribution and social behavior of some pelagic mysids shrimps. J. Exp. Mar. Biol. Ecol., 3, 125-155.
- Cochlan W.P., Wilkner J., Steward D.C., Smith D.C., Azam F., 1993 Spatial distribution of viruses, bacteria and chlorophyll *a* in neritic, oceanic and estuarine environments. Mar. Ecol. Prog. Ser., 92, 77-87.
- Codispodi L.A., 1989. Phosphorus vs. Nitrogen limitation of new and export production. In W.H. Berger, V.S. Smetacek & G. Wefer (Eds): Productivity of the ocean: present and past, Wiley & Sons, Chichester, pp 377-394.
- Cole J.J., Findlay S., Pace M.L., 1988. Bacterial production in fresh and salt water ecosystems: A cross-system overview. Mar. Ecol. Prog. Ser., 43, 1-10
- Collos Y., Slawyk G., 1980. Nitrogen uptake and assimilation by marine phytoplankton. In P.G. Falkowski (Ed): Primary Productivity in the Sea, Plenum Press, pp. 195-211.
- Comiso J.C., 1991. Satellite remote sensing of the polar oceans. J. Mar. Syst., 2, 395-434.
- Comiso J.C., McClain C.R., Sullivan C.W., Ryan J.P., Leonard C.L., 1993. Coastal zone color scanner pigment concentration in the Southern Ocean and relationships to geophysical surface features. J. Geophys. Res., 98, 2419-2451.
- Conover R.J., 1960. The feeding behavior and respiration of some marine planktonic crustacea. Biol. Bull., 119, 399-415.
- Conover R.J., 1966a. Assimilation of organic matter by zooplankton. Limnol. Oceanogr., 11, 338-345.
- Conover R.J., 1966b. Factors affecting the assimilation of organic matter by zooplankton and the question of superfluous feeding. Limnol. Oceanogr., 11, 346-354.
- Conover R.J., 1966c. Feeding on large particles by *Calanus hyperboreus* (Kröyer). In Barnes (Ed), Some contemporary studies in marine science. Allen & Unwin, Lond., pp. 187-194.
- Conover R.J., 1982. Interrelation between microzooplankton and other plankton organisms. Ann. Inst. Océanogr., Paris, 58(S), 31-46.
- Conover R.J., Corner E.D.S., 1968. Respiration and nitrogen excretion by some marine zooplankton in relation to their life cycle. J. Mar. Biol. Ass. U.K., 48, 49-75.
- Corner E.D.S., Newell B.S., 1967. On the nutrition and metabolism of zooplankton. IV. The forms of nitrogen excreted by *Calanus*. J. Mar. Biol. Ass. U.K., 47, 113-120.
- Coste B., Gostan J., Minas H.J., 1972. Influence des conditions hivernales sur les productions phyto- et zooplanctoniques en Méditerranée Nord-Occidentale. I. Structures hydrologiques et distribution des sels nutritifs. Mar. Biol., 16; 320-348.
- Coste B., Le Corre P., Minas H.J., Morin P., 1988. Les éléments nutritifs dans le bassin occidental de la Méditerranée. Bilan des échanges avec l'Océan Atlantique à Gibraltar. Oceanol. Acta, 9 (Spec. Issue), 87-94.
- Cottrell M., Suttle C., 1991. Wide-spread occurrence and clonal variation in viruses which cause lysis of a cosmopolitan, eukaryotic marine phytoplankton, *Micromonas pusilla*. Mar. Ecol. Prog. Ser., 78, 1-9.
- Creazzo S., 1992. Meteorological observations. In: Oceanographic campaign 1989-1990 data report part II. Ministry of the University and Scientific and Technological Research. Republica Italiana., pp. 63-78.
- Cullen J.J., Eppley R.W., 1981. Chlorophyll maximum layers of the southern California Bight and possible mechanisms of their formation and maintenance. Oceanol. Acta, 4, 23-32.
- Curl H., 1962. Analysis of carbon in marine plankton organisms. J. Mar. Res. 207, 181-188.
- Cushing, D.H., 1975. Marine Ecology and Fisheries. Cambridge University Press, Cambridge, UK. 278 pp
- Da Costa J.A., Martin J.M., Meybeck M., 1980. Les apports fluviaux dans les pays du bassin méditerranéen. L'eau Pure, 56, 1-25.
- Dagg M.J., Walser Jr.W.E., 1986. The effect of food concentration on fecal pellet size in marine copepods. Limnol. Oceanogr., 31, 1066-1071.

- Daro M.H., 1988. Migratory and grazing behaviour of copepods and vertical distribution of phytoplankton. Bull. Mar. Sc., 43, 710-729.
- Daro M.H., Hecq, J.H., 1983. Variations nyctémérales du broutage, de l'activité amylasique et du taux respiratoire du zooplancton en baie de Calvi (Corse). Rapp. Comm. Int. Mer Médit., 28, 227-228.
- Dauby P., 1980. Cycle annuel du zooplancton de surface de la baie de Calvi (Corse). Biomasse totale et plancton copépodien. Oceanol. Acta, 3, 403-407.
- Dauby P., 1985. Dynamique et productivité de l'écosystème planctonique du golfe de Calvi-Corse. Ph.D. Thesis, Univ. Liège, Belgium, 291 pp.
- Davidson A.T., Marchand H.J., 1992. The biology and ecology of *Phaeocystis* (Prymnophyceae). Progress in Physiological Research, 8, 1-45.
- Davis C.S., Flierl G.R., Wiebe P.H., Franks P.J.S., 1991. Micropatchiness, turbulence and recruitment in plankton. J. Mar. Res., 49, 109-151.
- Davis C.S., Gallager S.M., Solow A.R., 1992. Microaggregations of oceanic plankton observed by towed video microscopy. Science, 257, 230-232.
- Delhez E.J.M., Lacroix G., Martin G., Nihoul J.C.J., Varela R.A., 1993. Development of interdisciplinary models of ecosystems under severe hydrodynamic constraints. In: Progress in Belgian Oceanography Research (Brussels, January 21-22 1993), Royal Academy of Belgium, National Committee of Oceanology, pp 37-54.
- Dieckman G.G., Lange M.A., Ackley, Jennings Jr J.C., 1991. The nutrient status in sea ice of the Weddell Sea during winter: effects of sea ice texture and algae. Polar Biol., 11, 449-456.
- Djenidi S., 1985. Observations au large de Calvi (Corse) en régime d'été. Bull. Soc. Roy. Sci. Liège, 54, 287-300.
- Doney, S.C., Glover, D.M., Najjar, R.G., 1996. A new coupled one-dimensional biological-physical model for the upper ocean: applications to the JGOFS Bermuda Atlantic Time-series Study (BATS) site. Deep-Sea Res., 43, 591-624.
- Doronin Y.P., Kheisin D.E., 1977. Sea ice. Amerind, 323 pp.
- Ducklow H.W., Fasham M.J.R., 1992. Bacteria in the greenhouse: Modelling the role of oceanic plankton in the global carbon cycle. In R. Mitchell (Ed): Environmental Microbiology. Wiley, New York.
- Dugdale R.C., 1967. Nutrient limitation in the sea: dynamics, identification and significance. Limno. Oceanogr., 12, 685-695.
- Dugdale R.C., Goering J.J., 1967. Uptake of new and regenerated forms of nitrogen in primary productivity. Limnol. Oceanogr., 12, 179-184.
- Dugdale R.C., Mac Isaac J.J., 1971. A computation model for the uptake of nitrate in the Peru upwelling region. Invest. Pesq., 38, 299-308.
- Dugdale R.C., Wilkerson F.P., 1988. Nutrient sources and primary production in the Eastern Mediterranean. Oceanol. Acta, 9 (Spec. Issue), 179-184.
- Dugdale R.C., Wilkerson F.P., 1991. Low specific uptake rate: A common feature of high nutrient, low-chlorophyll marine ecosystems. Limnol. Oceanogr., 36, 1678-1688.
- Dunbar R.B., Anderson J.B., Domack E. W., Jacobs S.S., 1985. Oceanographic influences on sedimentation along the antarctic continental shelf. Oceanology of the Antarctic Continental Shelf. Antarctic Res. Ser., 13, 291-312.
- Dussart B.M., 1965. Les différentes catégories de plankton. Hydrobiologia, 26, 72-74.
- Eberlein K., Leal M.T., Hammer D.K., Hickel W., 1985. Dissolved organic substance during a *Phaeocystis pouchetii* bloom in the German Bight. Mar. Biol., 89, 311-316.
- Edmonson W.T., 1961. Food supply and reproduction of zooplankton in relation to phytoplankton populations. Rapp. P.-V. Réun. Cons. Perm. Int. Explor. Mer. 153, 137-141.
- Edmonson W.T., Winberg G.C., 1971. A manual on methods for the assessment of secondary productivity in freshwaters. I.B.P. Handbook n°17, Blackwell, Oxford and Edinburgh, 358pp.
- Elkalay K., Frangoulis C., Skliris N., Goffart A., Gobert S., Lepoint G. & Hecq JH (2003) A model of the seasonal dynamics of biomass and production of the seagrass *Posidonia oceanica* in the Bay of Calvi (Northwestern Mediterranean). *Ecol. Model.* 167: 1-18

- Elkhay K., N. Skliris, C. Frangoulis, A. Goffart, S. Gobert, G. Lepoint, J.H. Hecq, 2000. A model describing the *Posidonia oceanica* growth, *Biol. Mar. Medit.* 747-50
- Elliot A.M., Clemmons G.L., 1966. An ultrastructural study of ingestion and digestion in *Tetrahymena pyriformis*. *J. Protozool.*, 13, 311-323.
- El-Sayed S.Z., Holm-Hansen, O., Biggs D.C., 1983. Phytoplankton standing crop, primary productivity, and near surface nitrogenous nutrient fields of the Ross Sea, Antarctica. *Deep-Sea Research*, 30, 871-886.
- Engelkes M.M. 1985. Derivaatvorming en afbraak van algenpigmenten door copepodenbegrazing gemeten na scheiding met HPLC. Studentenverslag, N.I.O.Z., Texel, The Netherlands.
- Eppley R.W., 1972. Temperature and phytoplankton growth in the sea. *Fish. Bull. US*, 70, 1063-1085.
- Eppley R.W., 1977. The growth and culture of diatoms. In D. Werner (Ed): *The Biology of Diatoms*. Blackwell Scientific, Oxford, p 24-31.
- Eppley R.W., 1986. Short term variations in primary productivity. In J.D. Burton, P.G. Brewer, R. Chesselet (Eds): *Dynamic processes in the chemistry of the upper ocean*. Nato Conf. Series 17, 187-195.
- Eppley R.W., Peterson B.J., 1979. Particulate organic matter flux and planktonic new production in the deep ocean. *Nature*, 282, 677-680.
- Estrada M., 1985. Deep phytoplankton and chlorophyll maxima in the Western Mediterranean. In M. Moraitou-Apostolopoulou & V. Kiortis (Eds): *Mediterranean Marine Ecosystems*, Plenum Press, New York, London 1, 247-277.
- Estrada M., Vives F., Alcaraz M., 1985. Life and the productivity of the open ocean. In R. Margalef (Ed): *Western Mediterranean*, Pergamon Press, pp. 148-197.
- Estrada M., Margalef R., 1988. Supply of nutrients to the Mediterranean photic zone along a persistent front. *Oceanol. Acta*, 9 (Spec. Issue), 133-142.
- Evans G.T., Taylor F.J.R., 1980. Phytoplankton accumulation in Langmuir cells. *Limnol. Oceanogr.*, 25, 840-845.
- Everson I., 1982. Diurnal variations in mean volume back scattering strength of Antarctic krill (*Euphausia superba*) patch. *J. Plankton Res.*, 4, 155-162.
- Everson I., Bone D.G., 1986. Effectiveness of RMT8 system for sampling krill (*Euphausia superba*) swarms. *Polar Biol.*, 6, 83-90.
- Falkowski P.G., 1980. Light-shade adaptation in marine phytoplankton. In P. Falkowski (Ed): *Primary productivity in the sea*. Plenum Press, New York, London, pp 99-119.
- Falkowski P.G., Wirick C.D., 1981. A simulation model of the effects of vertical mixing on primary production. *Mar. Biol.*, 65, 69-75.
- Falkowski P.G., La Roche J., 1991. Adaptation to spectral irradiance in algae. *J. Phycol.*, 27, 8-14.
- Fasham M.J.R., Ducklow H.W., McKelvie S.M., 1990. A nitrogen-based model of plankton dynamics in the ocean mixed layer. *J. Mar. Res.*, 48, 591-639.
- Fasham M.J.R., Holligan P.P., Pugh P.R., 1983. The spatial and temporal development of the spring phytoplankton bloom in the Celtic Sea, April 1979. *Prog. Oceanogr.*, 12, 87-145.
- Fellous J.L., Morel A., 1987. French activities in ocean colour observations. In "Ocean Colour", European Space Agency, ESA SP-1083, 59-69.
- Fenaux R., 1967. Les appendiculaires des mers d'Europe et du Bassin Méditerranéen. *Faune Europe Bassin Méditerranéen*, 2, Masson, Paris, 166pp.
- Fenschel T., 1988. Marine plankton food chains. *Ann. Rev. Ecol. Syst.* 19, 19-38.
- Fenschel T., Blackburn, T.H., 1979. *Bacteria and mineral cycling*. Academic Press, London, 225pp.
- Ferrier C., Rassoulzadegan F., 1991. Density-dependant effects of protozoan on specific growth rates in pico- and nanoplanktonic assemblages. *Limnol. Oceanogr.*, 36, 657-669.
- Fichefet T., Gaspar P., 1988. A model study of upper ocean-sea ice interactions. *J. Phys. Oceanogr.*, 18, 181-195.
- Fleming R.H., 1939. The control of diatom populations by grazing. *J. Cons. Perm. Expl. Mer.*, 14, 210-227.

- Flos J. 1985. The driving machine. In R. Margalef (Ed): Western Mediterranean, Pergamon Press, pp. 61-99.
- Fonda Umani S., Monti M., Nuccio C., 1998. Microzooplankton biomass distribution in Terra Nova Bay, Ross Sea (Antarctica). In J. Le Fevre and P. Tréguer (Eds): Carbon Fluxes and Dynamic Processes in the Southern Ocean: Present and Past. J. Mar. Syst., 17, 289-303.
- Fortier L., Le Fèvre J., Legendre L., 1994. Export of biogenic carbon to fish and to deep ocean: the role of large planktonic microphages. J. Plankton Res., 8, 365-383.
- Fowler S.W., Fisher N.S., 1983. Viability of marine phytoplankton fecal pellets. J. Mar. Res., 30, 963-969.
- Foyn L., Rey F. 1981. Nutrient distribution along the Norwegien coastal current. In: The Norwegien Coastal Current, University of Bergen, 2, 629-648.
- Frangoulis C., Belkhiria S., Goffart A., Hecq J.H., 2001. Dynamics of copepod faecal pellets in relation to a *Phaeocystis* dominated bloom: characteristics, production and flux. J. Plankton Res., 23, 75-88.
- Franks P.J.S., Wroblewski J.S., Flierl G.R., 1986. Behavior of a simple plankton model with food-level acclimation by herbivores. Mar. Biol., 91, 121-129.
- Frontier S., 1985. Les fronts hydrologique: écosystèmes de contact. Dans Aspects interdisciplinaires de l'océanographie Méditerranée dans le domaine pélagique. Abstracts du Colloque international du CNRS, Villefranche-sur-Mer, 16-20/09/1985.
- Frontier S., 1986. Studying fronts as contact ecosystems. In In J.C.J. Nihoul (Ed): Marine Interfaces Ecohydrodynamics. Elsevier Oceanography Series, Amsterdam, 42, 55-66.
- Frontier S., 1999. Les écosystèmes. Presses Universitaires de France, 127 pp.
- Frost B.W., 1987. Grazing control of phytoplankton stock in the open subarctic Pacific Ocean: a model assessing the role of mesozooplankton, particularly the large calanoid copepods *Neocalanus* ssp. Mar Ecol. Progr. Ser., 39, 49-68.
- Frost B.W., 1991. The role of grazing in nutrient-rich areas of the open sea. Limnol. Oceanogr., 36, 1616-1630.
- Fuhrman J. A., Sleeter T.D., Carlson C.A., Proctor L.M., 1989. Dominance of bacterial biomass in the Sargasso sea and its ecological implications. Mar Ecol. Prog. Ser., 57, 207-217.
- Fuhrman J.A., Suttle C.A., 1993. Viruses in marine planktonic systems. Oceanography, 6, 51-63.
- Fukuchi M., Hattori H., Sasaki H., Hoshiai T. 1988. A phytoplankton bloom and associated processes observed with a long-term moored system in antarctic waters. Mar. Ecol. Prog. Ser., 45, 279-288.
- Garrison D.L., 1984. Planktonic diatoms. In K.A. Steidinger & L.M. Walker (Eds): Marine Plankton Life Cycle Strategies, CRC Press Boca Raton, Fla, pp 1-17
- Garrison D.L., Buck K.R., Fryxell G., 1987. Algal assemblages in Antarctic pack ice and in ice-edge plankton. J. Phycol., 23, 564-572.
- Gascard J.C., 1978. Mediterranean deep water formation. Baroclinic instability and oceanic eddies, Oceanologica Acta, 4, 315-330.
- Geider R.J., 1987. Light and temperature dependance of the carbon to chlorophyll ratio in microalgae and cyanobacteria: implication for physiology and growth of phytoplankton. New Phytol., 106, 1-34.
- Geider R.J., MacIntyre H.L., Kana T.M., 1997. A dynamic model of phytoplankton growth and acclimatation: responses of the balanced growth rate and the chlorophyll *a*: carbon ratio to light, nutrient-limitation and temperature. Mar. Ecol. Prog. Ser. 148, 187-200.
- George R.Y., 1984. Ontogenetic adaptationns in growth and respiration of *Euphausia superba* in relation to temperature and pressure. J. Crustac. Biol., 4, 252-262.
- Gieskes W.W.C., Kraay G.W., 1986. Analysis of phytoplankton by HPLC before, during and after mass occurrence of the microflagellate *Corymbellus aureus* during the spring bloom in the open northern North Sea in 1983. Mar. Biol. 92, 45-52.
- Gieskes W.W.C., Kraay G.W., Nontji A., Setiapermana D., Sutomo, 1988. Monsoonal alteration of a mixed and a layered structure in the phytoplankton of the euphotic zone of the Banda Sea (Indonesia): a mathematical analysis of algal pigments fingerprints. Neth. J. Sea Res., 22, 123-137.

- Glibert P.M., Biggs D.C., McCarthy J.J., 1982. Utilization of ammonium and nitrate during austral summer in the Scotia Sea. Deep Sea Res., 29, 837-850.
- GLOBEC, 1988. Global Ocean Ecosystems Dynamics. In V. Cullen (Ed): Report of a workshop held at Wintergreen, Virginia, May 1988, Joint Oceanographic Institutions, Inc. Publ., Washington, USA, 131 pp.
- Govert S., Belkhiria S., Dauby P., Havelange S., Soullard M., Bouquegneau J.M., 1995. Variations temporelles de la phénologie et de la composition biochimique de la phanérogamie du bassin marine *Posidonia oceanica* en baie de Calvi. Bull. Soc. Roy Sc. Liège, 64, 263-284.
- Godeaux J., 1988. Thalassocénés méditerranéens: une synthèse. Bull. Soc. Roy. Sci. Liège, 57, 359-377.
- Goffart A., 1992. Influence des contraintes hydrodynamiques sur la structure des communautés phytoplanctoniques du bassin Liguro-Provençal (secteur Corse). Ph.D. Thesis, Univ. Liège, 163 pp.
- Goffart A., Catalano G., Hecq J.H., 2000. Factors controlling the distribution of diatoms and *Phaeocystis* in the Ross Sea. J. Mar. Syst., 27, 161-175.
- Goffart A., Catalano G., Magazzù G., Hecq J.H., 1992. Some examples of the influence of hydrodynamical constraints on the phytoplanktonic biomass distribution in the Southern Ocean. In V. Gallardo, O. Feretti & H. Moyano (Eds): Acts from the International Seminar on the Oceanography in Antarctica, Concepcion, Chile (March 7-9 1991), pp. 265-271.
- Goffart A., Hecq J.H., 1985. Influence du front Liguro-provençal -secteur corse- sur la production planctonique. Rapp. Comm. Int. Mer Médit., 29, 133-134.
- Goffart A., Hecq J.H., 1987. Campagne INDIGO III 1987. Distribution spatiale du phyto- et du zooplankton en relation avec les masses d'eau. Résultats préliminaires. Programme ANTAR 06, Preliminary reports, 18 pp.
- Goffart A., Hecq J.H., 1988. Distribution of phytoplanktonic parameters in the Indian sector of the Southern Ocean during Indigo III cruise. In: Proceedings of the Belgian National Colloquium on Antarctic Research, Brussels, October 20, 1987, Prime Ministers Service Science Policy Office, pp. 147-166.
- Goffart A., Hecq J.H., 1989. Zooplankton biochemistry and ecodynamic. In S. Caschetto (Ed): Belgian Scientific Research Programme on Antarctica, Belgian Prime Minister's Services Sciences Policy Office. Brussels Belgium, 1, 1-60.
- Goffart A., Hecq J.H., 1990a. Stabilité interannuelle de la distribution de la production planctonique associée au front Liguro-Provençal (secteur Corse). Rapp. Comm. Int. Mer Médit., 32, 113.
- Goffart A., Hecq J.H., 1990b. Ecohydrodynamical approach of planktonic ecosystem in the Southern Ocean. Abstracts of "Symposium of the Biogeochemistry and the Circulation of the Southern Ocean" (Brest, France, 01-07 July 1990).
- Goffart A., Hecq J.H., 1993a. Biochemistry and ecodynamics of zooplankton of the Southern Ocean. In S. Caschetto (Ed): Belgian Scientific Research Programme on the Antarctic, Scientific Results of phase two, 56 pp.
- Goffart A., Hecq J.H., 1993b. Control of phytoplankton development by nitrate availability in the Liguro-provençal basin (Western Mediterranean). In: Progress in Belgian Oceanography Research (Brussels, January 21-22 1993), Royal Academy of Belgium, National Committee of Oceanology, pp 251-261.
- Goffart A., Catalano G., Hecq J.H., 2000. Factors controlling the distribution of diatoms and *Phaeocystis* in the Ross Sea., Journal of Marine Systems, 27, 161-175.
- Goffart A., Hecq J.H., Licot M., 1985. Ecohydrodynamical study of the Liguro-Provençal Front (Corsica). II. Phytoplankton Data. In R. Van Grieken & R. Wollast (Eds): Proceedings "Progress in Belgian Oceanographic Research", Brussels, March 1985, University of Antwerp, pp. 408-413.
- Goffart A., Hecq J.H., Norro A., Petit B., Wilmette A., 1999. Southern Ocean global ecosystem response to physical and trophic constraints. Mid-term report. Programme "Recherches Scientifiques sur l'Antarctique - phase IV". Services du Premier Ministre, Services Fédéraux des Affaires Scientifiques, Techniques et Culturelles.

- Goffart A., Hecq J.H., Prieur, L. 1995. Contrôle du phytoplancton du Bassin Ligur par le Front Liguro-Provençal (secteur corse). *Oceanol. Acta*, 18, 329-342.
- Goffart A., Licot M., Mouchet A., Hecq J.H., 1988. Le système frontal Liguro-Provençal et ses implications biologiques au large de Calvi. *Bull. Soc. r. Sc. Liège*, 17, 346.
- Goldman J.C., 1980. Physiological processes, nutrient availability, and the concept of relative growth rate in marine phytoplankton ecology. In P.G. Falkowski (Ed): Primary productivity in the sea. Plenum Press, New York, 179-193.
- Goldman J.C., 1984. Oceanic nutrient cycles. In M.J.R. Fasham (Ed): Flows of energy and materials in marine ecosystems, Nato Conf. IV Marine Sciences Series, Plenum. New York, 13, 137-170.
- Goldman J.C., 1988. Spatial and temporal discontinuities of biological processes in pelagic surface waters. In B.J. Rothschild (Ed): Toward a Theory on Biological-Physical Interactions in the World Ocean, Kluwer Academic Publisher, pp. 273-296.
- Goldman J.C., 1993. Potential role of large oceanic diatoms in new primary production. *Deep-Sea Res.*, 40, 159-168.
- Goldman J.C., Caron D.A., Dennett M.R., 1987. Regulation of gross growth efficiency and ammonium regeneration in bacteria by substrate C:N ratio. *Limnol. Oceanogr.*, 32, 1239-1252.
- Goeyens L., Elskens M., Catalano G., Lipizer M., Hecq J.H., Goffart A., 2000. Nutrient depletions in the Ross Sea and their relation with pigment stocks. *J. Mar. Syst.*, 27, 195-208.
- Goosse H., Hecq J.H., 1994. Modelling the ice-ocean-plankton interactions in the Southern Ocean. *J. Mar. Syst.*, 5: 471-484.
- Grégoire M.L., Beckers J.M., Nihoul J.C.J., Stanev E., 1998. Reconnaissance of the Black Sea' ecohydrodynamics by means of a 3D interdisciplinary model. *J. Mar. Syst.*, 16, 85-105.
- Guglielmo L., Hecq J.H., 1990. Preliminary results of the Vth ITALIANTARTIDE Oceanographic Expedition in Ross Sea and in the Pacific sector of the Southern Ocean. Abstracts of the 22nd International Liège Colloquium on Ocean Hydrodynamics, pp. 28-29.
- Guillard R.L., Killam P., 1977. The ecology of marine planktonic Diatoms. In O. Kinne (Ed): Marine Ecology. IV. Dynamics, Wiley & Sons, Chichester, pp 372-469.
- Guillard R.R.L., Murphy L.S., Foss P., Liaaen-Jensen S., 1985. *Synechococcus* spp. as likely zeaxanthin-dominant ultraphytoplankton in the North Atlantic. *Limnol. Oceanogr.*, 30, 412-414.
- Harding L.W., 1988. The time-course of photoadaptation to low-light in *Prorocentrum marielebouriae* (Dinophyceae). *J. Phycol.*, 24, 274-281.
- Hargraves P.E., 1976. Studies on marine planktonic diatoms. II resting spores morphology. *J. Phycol.*, 12, 118-132.
- Harris E., 1959. The nitrogen cycle in Long Island Sound. *Bull. Bing. Oceanogr. Coll.*, 17, 31-65.
- Harris G.P., 1980. The measurement of photosynthesis in natural populations of phytoplankton. In I. Morris (Ed): The physiological ecology of phytoplankton, Blackwell Scientific Publications, pp. 129-187.
- Harrison W.G., Wood L.J.E., 1988. Inorganic nitrogen uptake by marine picoplankton: evidence for size partitioning. *Limnol. Oceanogr.*, 33, 468-475.
- Haury L.R., McGowan J.A., Wiebe P.H., 1978. Patterns and processes in the time-space scales of plankton distributions. In J.H. Steele (Ed): Spatial pattern in plankton communities, Plenum Press, New York, pp. 277-327.
- Hay S.J., Kiorboe T., Matthews A., 1991. Zooplankton biomass and production in the North Sea during the Autumn Circulation Experiment, October 1987 - Mars 1988.
- Hecq J.H., 1975. Cycle annuel du zooplancton à Boulogne sur mer (Pas de Calais). I. Le Méroplancton. *Bull. soc. r. sci. Liège*, 44, 477-485.
- Hecq J.H., 1976. Cycle annuel du zooplancton à Boulogne sur mer (Pas de Calais). II. L'Holoplancton. *Bull. Soc. r. Sc. Liège*, 45, 443-447.
- Hecq J.H., 1979a. Rôle du zooplancton dans la cycle des nutriments. Application au Pas de Calais et au sud de la mer du Nord. *Rapp. Comm. Int. Mer Médit.*, 25-26, 109.
- Hecq J.H., 1979b. Caractérisation des masses d'eau de petite dimension par les indicateurs zooplanctoniques. *Rapp. Comm. Int. Mer Médit.*, 25-26, 117-118.

- Hecq J.H., 1982. Distribution et dynamique du zooplancton en relation avec le cycle du carbone en baie sud de la mer du Nord. PhD Thesis. University of Liege, 182 pp.
- Hecq J.H., Bouquegneau J.M., Djenidi S., Frankignoulle M., Goffart A., Licot M., 1986. Some aspect of the Liguro-provençal frontal ecohydrodynamics. In J.C.J. Nihoul (Ed): Marine Interfaces Ecohydrodynamics. Elsevier Oceanography Series, Amsterdam, 42, 257-271.
- Hecq J.H., Brasseur P., Goffart A., Lacroix G., Guglielmo L., 1993. Modelling approach of the planktonic vertical structure in deep Austral Ocean. The example of the Ross Sea Ecosystem. In: Progress in Belgian Oceanographic Research. Proceedings of the annual workshop on Belgian Oceanographic Research, 235-250.
- Hecq J.H., Gaspar A., 1980. Composition biochimique du zooplancton en mer du Nord. Approche écologique. Ann. Soc. R. zool. de Belgique, 110, 105-113.
- Hecq J.H., Gaspar A., Dauby P., 1981a. Caractéristiques écologiques et biochimiques de l'écosystème planctonique en baie de Calvi. Premières Journées d'Océanographie Méditerranéennes. Bull. Soc. r. Sc. Liège, 50, 440-445.
- Hecq J.H., Gaspar A., Dauby P., 1981b. Variation annuelle de la composition biochimique du zooplancton en baie de Calvi (Corse). Rapp. Comm. Int. Mer Médit., 27, 203-204.
- Hecq J.H., Goffart A., 1984. Analyse des classes de lipides et des acides gras de *Leptomyysis lingvura* (Sars), crustacés mysidacé. Influence des conditions nutritionnelles. Bull. Soc. r. Sc. Liège, 53, 69-80.
- Hecq J.H., Goffart A., 1990a. Influence of frontal systems on planktonic communities in Austral Ocean. Abstracts of the 22nd International Liège Colloquium on Ocean Hydrodynamics, 31-32.
- Hecq J.H., Goffart A., 1990b. Le système frontal Liguro-Provençal et ses implications biologiques sur le plateau continental Corse au large de Calvi. Programme "Flux Océaniques" (France - JGOFS), Frontal, Rapp.n°8, 104-107.
- Hecq J.H., Goffart A., 1996. The pivotal role of zooplankton in the North Sea pelagic ecosystem. In OSTC (Ed.): Dialog between scientists and users of the Sea. Proceedings of a symposium held on the occasion of the 10th anniversary of the civil service oceanographic research vessel Belgica, Ostend, 17 - 19 October 1994, pp. 21-32.
- Hecq J.H., Goffart A., Catalano G., 1991. An ecosystem approach of the Straits of Magellan, based on nutrients concentrations and phytopigments characteristics. Memorie di Biologia Marina e di Oceanografia, 19, 97-99.
- Hecq J.H., Goffart A., Licot M., Dauby P., Brohée M., 1988. L'écosystème planctonique en baie de Calvi (Corse): Caractéristiques biochimiques et physiologiques. Soc. r. Sc. Liège, 17, 347.
- Hecq J.H., Goffart A., Nihoul J.C.J., 1986. Plankton dynamic at marine frontal interfaces. Eos Transaction. American Geophysical Union, 67, 982.
- Hecq J.H., Guglielmo L., Goffart A., Catalano G., Goosse H., 1999. Modeling approach of the Ross Sea plankton ecosystem. In F. Faranda, L. Guglielmo & A. Ianora (Eds): Ross Sea Ecology. Springer, Berlin Heidelberg, pp. 395-411.
- Hecq J.H., Guglielmo L., 1992. Structure and functioning of the Ross Sea pelagic ecosystem: an Interdisciplinary approach. In V. Gallardo, O. Feretti & H. Moyano (Eds): Acts from the International Seminar on the Oceanography in Antarctica, Concepcion, Chile (March 7-9 1991), pp. 227-233.
- Hecq J.H., Heiden D., Moermans R., 1976. Biologie de *Sagitta setosa* dans le Pas de Calais et le sud de la mer du Nord. Bull. Soc. r. Sc. Liège, 45, 653-667.
- Hecq J.H., Licot M., Goffart A., Mouchet A., Frankignoulle M., Bouquegneau J.M., Distèche A., Godeaux J., Nihoul J.C.J., 1985. Ecohydrodynamical study of Liguro-Provençal front (Corsica). I. Hydrological Data. In R. Van Grieken & R. Wollast (Eds): Proceedings "Progress in Belgian Oceanographic Research", Brussels, March 1985. University of Antwerp, pp. 397-407.
- Hecq J.H., Marty J.C., Claustre H., Prieur L., 1992a. Distributions des produits de dégradation de la chlorophylle *a* durant la campagne Almofront-1. In A. Sournia & L. Prieur (Eds): Processes and budgets in geostrophic fronts (France JGOFS), pp. 116-118.

- Hecq J.H., Magazzù G., Goffart A., Catalano G., Vanucci S., Guglielmo L., 1992b. Distribution of planktonic components related to vertical structure of water masses in the Ross Sea and the Pacific sector of the Southern Ocean. Atti del 9° congresso A.I.O.L., Santa Margherita Ligure, 20-23 novembre 1990, pp. 665-678.
- Hecq J.H., Mingelbier M., Goffart A., Brylinsky J.M., Djenidi S., 1992c. Caractéristiques écohydrodynamiques de la baie sud de la mer du Nord en régime d'été. Bull. Soc. r. Sc. Liège, 61, 79-97.
- Hecq J.H., Sainvitu E., Dauby P., 1984. Influence des conditions expérimentales et environnementales sur la consommation d'oxygène et l'activité nutritionnelle de *Leptomyysis lingvura* (Sars) (Crustacé, Mysidacé). Annls. Soc. r. Zool. Belg., 114, 200-201.
- Hecq J.H., Norro A., Goffart A. 2001. Modelling the influence of ice coverage variability on the Ross Sea Plankton Assemblages. Submitted to Journal of Marine Systems
- Herman A.W., Samaeoto D.D., Longhurst A.R., 1981. Vertical and horizontal distribution patterns of copepods near the shelf break South of Nova Scotia. Can. J. Fish. Aquat. Sci., 38, 1065-1076.
- Hirche H.J., 1983. Overwintering of Calanus finmarchicus and Calanus helgolandicus. Mar. Ecol. Prog. Ser., 11, 281-290.
- Hofmann E.E., Ambler J.W., 1988. Plankton dynamics on the outer southeastern U.S. continental shelf. Part II: A time-dependent biological model. J. Mar. Res., 46, 883-917.
- Holligan P.M., Williams P.J., Purdie D., Harris R.P., 1984. Photosynthesis, respiration and nitrogen supply of plankton populations in stratified, frontal and tidally mixed shelf waters. Mar. Ecol. Prog. Ser., 17, 201-213.
- Holligan P.M., Aarup T., Groom S.B., 1989. The North Sea: Satellite Colour Atlas. Cont. Shelf Res., 9(8), 667-765.
- Holm-Hansen O., 1985. Nutrient cycles in Antarctic marine ecosystems. In W.R. Siegfried, P.R. Condy and R.M. Laws (Eds). Antarctic Nutrient Cycles and Food Webs. Springer-Verlag, Berlin, p 6-10.
- Honjo S., 1976. Coccoliths: Production transportation and sedimentation. Mar. Micropaleont. 1, 65-79.
- Honjo S., Emery K.O, 1976. Suspended matter of eastern Asia: Scanning electron microscopy and X-Ray peak analysis. In H. Aoki, S. Iisuka (Eds): Volcanoes and technosphère. Tokai Univ. Japan, pp259-288.
- Honjo S., Roman M.R., 1978. Marine copepod fecal pellets: production preservation and sedimentation. J. Mar. Res. 36, 45-57.
- Hopkins C.C.E., Tande K.S., Gronvik S., Sargent J.R., 1984. Ecological investigations on the zooplankton community of Balsfjorden, northern Norway: an analysis of growth and overwintering tactics in relation to niches and environment in *Metridia longa* (Lubbock), *Calanus finmarchicus* (Gunnerus), *Thysanoessa inermis* (Kroyer) and *T. raschi* (Sars). J. Exp. Mar. Biol. Ecol., 88, 77-89.
- Hudson R.J.M., Morel F.M.M., 1990. Iron tranport in marine phytoplankton: Kinetics of cellular and medium coordination reactions. Limnol. Oceanogr. 35, 1002-1020.
- Hutchinson G.E., 1967. A treatise on Limnology, Vol. 2. Wiley, New York; 1116 pp.
- Ianora A., Mazzocchi M.G., Grottoli R., 1992. Seasonal fluctuations in fecundity and hatching success in the planktonic copepod *Centropages typicus*. J. Plankton Res., 14, 1483-1494.
- Ianora A., Poulet S.A., 1993. Egg viability in the copepod *Temora stylifera*. Limnol. Oceanogr., 38, 1615-1626.
- Ianora A., Poulet S.A., Miraldo A., 1995. A comparative study of the inhibitory effect of diatoms on the reproductive biology of the copepod *Temora stylifera*. Mar. Biol., 121, 533-539.
- Ibarra S.O., 1981. Contributions relatives du micro- et du nanophytoplancton à la production primaire dans une zone côtière de Méditerranée nord-occidentale. Vie et Milieu, 31, 119-128.
- Ichimura S.S., 1956. On the ecological meaning of transparency for the production of matter in phytoplankton community of lake. Bot. Mag., Tokyo, 69, 219-226.
- Idso S. B., Jackson R.D., 1969. Thermal radiation from the athmosphere. J. Geophys. Res., 74, 5397-5403.

- Ivanoff A., Morel A., 1970. Terminologie concernant l'optique océanographique. Cah. Océanogr., 22, 457-468.
- Ivlev V. S., 1945. The biological productivity of waters. Usp. Sovrem. Biol., 131-150.
- Ivlev V.S., 1955. Experimental ecology of the feeding of fishes. Yale Univ. Press, New Haven, 302 pp.
- Iizumi H., Akihiko H., 1982. Growth and organic production of eelgrass (*Zostera marina* L.) in temperature waters of the Pacific coast of Japan. III. The kinetics of nitrogen uptake. Aquat. Bot., 12, 245.-252.
- Jacobs S.S., Comiso J.C. 1989. Sea ice and oceanic processes on the Ross Sea continental shelf. J. Geophys. Res., 94, 18195-18211.
- Jacobsen D.M., 1987. The ecology and feeding biology of thecate dinoflagellates. Ph. D. Thesis WHOI-87-10.
- Jacques G., 1970. Aspects quantitatifs du phytoplancton de Banuyls-sur-Mer (Golfe du Lion). IV. Biomasse et production, 1965-1969. Vie et Milieu, 21 B, 37-102.
- Jacques G., 1988. Flux de carbone en milieu pélagique de Méditerranée Occidentale lors de la floraison printanière. Oceanol. Acta, 9 (Spec. Issue), 143-148.
- Jacques G., Tréguer P., 1986. Ecosystèmes pélagiques marins. Collection d'écologie, Masson, 243 pp.
- Jamart B.M., Winter D.F., Banse K., Anderson G.C., Lam R.K., 1977. A theoretical study of phytoplankton growth and nutrient distribution in the Pacific Ocean off the northwestern U.S. coast. Deep-Sea Res., 24, 753-773.
- Jamart B.M., Winter D.F., Banse,K., 1979. Sensitivity analysis of a mathematical model of phytoplankton growth and nutrient distribution in the Pacific Ocean off the northwestern U.S. coast. J. Plankton Res., 1, 267-290.
- Jassby A.D., Platt T., 1976. Mathematical formulation of the relationship between photosynthesis and light for phytoplankton. Limnol. Oceanogr., 21, 540-546.
- Jawed M., 1969. Body nitrogen and nitrogenous excretion in *Neomysis rayii* Murdoch and *Euphausia pacifica* Hansen. Limnol. Oceanogr., 14, 748-754.
- Jawed M., 1973. Ammonia excretion by zooplankton and its significance to primary productivity during summer. Mar. Biol., 23, 115-120.
- Jeffrey S.W., 1974. Profiles of photosynthetic pigments in the ocean using thin-layer chromatography. Mar. Biol., 26, 101-110.
- Jeffrey S.W., 1980. Algal pigment systems. In P. Falkowski (Ed): Primary productivity in the sea. Plenum Press, New York, pp. 35-58.
- Jennings J.C., Gordon L.I., Nelson D.M., 1984. Nutrient depletion indicates high primary productivity in the Weddell Sea. Nature, 309, 51-54.
- Jenssen A., Sakshaug E., 1973. Studies on the phytoplankton ecology of the Trondheimsfjord. II. Chloroplast pigments in relation to abundance and physiological state of the phytoplankton. J. Exp. Mar. Biol. Ecol., 11, 137-143
- Joint I.R., Pomroy A.J., 1987. Activity of heterotrophic bacteria in the euphotic zone of the Celtic Sea. Mar. Ecol. Prog. Ser., 41, 155-165.
- Joinis C., Billen G., Lancelot C., Mommaerts J.P., Daro M.H., Bossicart M., Gillain G., Bertels A., Hecq J.H., Wijnant P., 1979. Dynamic of organic matter in three planctonic ecosystems. Int. Counc. Explor. Sea. Oceanograph. commitee, 33, 165-184.
- Joinis C., Billen G., Lancelot C., Daro M.H., Mommaerts J.P., Bertels A., Bossicart M., Nijs J., Hecq J.H., 1982. A budget of carbon cycling in the Belgian coastal zone: relative roles of zooplankton, bacterioplankton and benthos in the utlization of primary production. Neth. J. Sea Res., 16, 260-275.
- Jorgensen S.E., 1988. Fundamentals of ecological modelling. developpements in environmental modelling. Elsevier, 391 pp.
- Jorgensen S.E., Nielsen S.N., Jorgensen L.A., 1991. Handbook of ecological parameters and ecotoxicology. Elsevier, 1263 pp.

- Jumars P.A., Perry D.L., Baross J.A., Perry M.J., Frost B.W., 1989. Closing the microbial loop: Dissolved carbon pathway to heterotrophic bacteria from complete ingestion, digestion and absorption in animals. Deep-Sea Res., 36, 483-495.
- Kirchman D.L., Lancelot C., Fasham M., Legendre L., Radach G., Scott M., 1993. Dissolved organic matter in biogeochemical models of the oceans. In G.T. Evans and M.J.R. Fasham (Eds): Towards a model of ocean biogeochemical processes. Springer-Verlag, Berlin. pp209-225.
- Klein B., Gieskes W.C., Kraay G.G., 1986. Digestion of chlorophylls and carotenoids by the marine protozoan *Oxyrrhis marina* studied by HPLC. analysis of algal pigments. J. Plankton Res., 8, 827-836.
- Klein B., Sournia A., 1987. A daily study of the diatom spring bloom at Roscoff (France) in 1985. II. Phytoplankton pigments composition studied by HPLC analysis. Mar. Ecol. Prog. Ser., 37, 265-275.
- Knox G.A., 1970. Antarctic marine ecosystems. In M.W. Holdgate (Ed): Antarctic Ecology. London, Academic Press, 1, 69-96.
- Knox G.A., 1994. The biology of the Southern Ocean. Cambridge University Press, 444 pp.
- Koehl M.A.R., Strickler J.R., 1981 Copepods feeding currents: food capture at low reynolds number. Limnol. Oceanogr. 26, 1062-1073.
- Koopmann G., 1953. Entstehung und Verbreitung von Divergenzen in der oberflächennahen Wasserbewegung der antarktischen Gewässer. Deutsche hydrographische Zeitschrift, Ergänzungsheft, 2, 1-17.
- Krause M., Martens P., 1990. Distribution patterns of mesoplankton biomass in the North Sea. Helgoländer Meeresunters., 44, 295-327.
- Krause M., Trahms J., 1982. Vertical distribution of copepods (all developmental stages) and other zooplankton during spring bloom in the Fladden Ground area of the North Sea. Neth. J. of Sea Res., 16, 217-230.
- Kremer J.N., Nixon S.W., 1978. A coastal marine ecosystem. simulation and analysis. Ecological Studies. Springer Verlag, Heidelberg, 24, 217 pp.
- Kumar S.K., Vincent W.F., Austin P.C., Wake G.C., 1991. Picoplankton and marine food chain dynamics in a variable mixed layer: a reaction-diffusion model. Ecol. Modelling, 57, 193-219.
- Kuramoto S., Koyama K., 1987. Preliminary report of the oceanographic observations on the 22nd Japanese Antarctic Research Expedition (1980-1981). Memoirs National Institute of Polar Research, Special Issue, 23, 5-12.
- Lacroix G., 1998. Simulation de l'écosystème pélagique de la mer Ligure à l'aide d'un modèle unidimensionnel. Etude du bilan de matière et de la variabilité saisonnière, interannuelle et spatiale. Ph.D. Thesis, Univ. Liège, Belgium and Univ. Pierre and Marie Curie, Paris VI, France, 256 pp.
- Lacroix G., Nival, P. 1999. Influence of meteorological variability on primary production dynamics in Ligurian Sea (NW Mediterranean Sea) with a 1D hydrodynamic/biological model. J. Mar. Syst., 16, 23-50.
- Lancelot C., Mathot S., Owens N.J.P., 1986. Modelling protein synthesis, a step to an accurate estimate of net primary production: *Phaeocystis pouchetii* colonies in Belgian coastal waters. Mar. Ecol. Prog. Series, 32, 193-202.
- Lancelot C., Veth C. and Mathot S., 1991. Modelling of the ice edge phytoplankton bloom in the Scotia Weddell Sea sector of the Southern Ocean. J. Mar. Syst., 2, 333-346.
- Lancelot C., Mathot S., Veth C., de Baar H., 1993. Factors controlling phytoplankton ice-edge blooms in the marginal ice-zone of the northwestern Weddell Sea during sea ice retreat 1988: field observations and mathematical modelling. Polar Biol., 13, 377-387.
- Lancelot C., Hannon E., Becquevort S., Veth C., De Baar H.J.W., 2000. Modeling phytoplankton blooms and carbon export production in the Southern Ocean: dominant controls by light and iron in the Atlantic sector in Austral spring 1992. Deep-Sea Research, I 47, 1621-1662.
- Langdon C., 1988a. On the cause of interspecific difference in the growth-irradiance relationship for phytoplankton. I. A comparative study of the growth irradiance relationship of three marine

- phytoplankton species: *Skeletonema costatum*, *Olisthodiscus luteus* and *Gonyaulax tamarensis*. J. Plankton Res., 10, 1291-1312.
- Lasker R., 1966. Feeding, growth, respiration and carbon utilization of an euphausiid crustacean. J. Fish. Res. Bd. Canada, 23, 1291-1317.
- LeBraseur R.J., Fulton, J., 1967. A guide to zooplankton of the north western Pacific Ocean. Fish Res. Bd. Canada, Circular, 84, 34pp.
- Lee A.J., 1970. The currents and water masses of the North Sea. Oceanogr. Mar. Biol., 8, 33-71.
- Le Fevre J., Frontier S., 1988. Influence of temporal characteristics of physical phenomena on plankton dynamics, as shown by northwest European marine ecosystems. In B.J. Rothschild (Ed.) Towards a Theory on Biological-Physical Interactions in the World Ocean. Kluwer, Dordrecht. Pp 245-272.
- Le Feuvre J., Legendre L., Rivkin R.B., 1998. Fluxes of biogenic carbon in the Southern Ocean: roles of large microphagous zooplankton. In J. Le Feuvre and P. Tréguer (Eds): Carbon Fluxes and Dynamic Processes in the Southern Ocean: Present and Past. J. Mar. Syst., 17, 325-345.
- Legendre L., 1981. Hydrodynamic control of marine phytoplankton production: the paradox of stability. In J.C.J. (Ed.) Marine Interfaces Ecohydrodynamics. Elsevier Oceanogr., Amsterdam. pp. 191-206.
- Legendre L., 1990. The significance of microalgal blooms for fisheries and for the export of particulate organic carbon in oceans. J. Plankton Res., 12, 681-699.
- Legendre L., Ackley S.F., Dieckmann G.S., Gulliksen B., Horner R., Hoshiai T., Melnikov I.A., Reeburgh W.S., Spindler M., Sullivan C.W., 1992. Ecology of sea ice biota, 2. Global significance. Polar Biol., 12, 429-444.
- Legendre L., Gosselin M., 1989. New production and export of organic matter to the deep ocean: consequence of some recent discoveries. Limnol. Oceanogr., 34, 1374-1380.
- Legendre L., Le Feuvre J., 1995. Microbial food webs and the export of biogenic carbon in oceans. Aquat. microb. Ecol., 9, 69-77.
- Legendre L., Rassoulzadegan F., 1995. Plankton and nutrient dynamics in marine waters. Ophelia, 41, 153-172.
- Lessard E.J., Smith J.R., 1986. Dinoflagellates from the North Atlantic classified as phototrophic or heterotrophic by epifluorescence microscopy. Journ. Plankt. Res., 8, 1209-1215.
- Levitus S. 1982. Climatological atlas of the world ocean. NOAA Prof , US Govt Print Off, Washington, DC. Pap 13, 1-173.
- Lewis J.B., 1981. Coral reef ecosystems. In A.R. Longhurst (Editor), Analysis of marine ecosystems. Academic press, 127-159.
- Lewis M.R., Cullen J.J., Platt T., 1983. Phytoplankton and thermal structure in the upper ocean: Consequences of nonuniformity in chlorophyll profile. J. Geophys. Res., 88, 2565-2570.
- Lewis MR., Harrison W.G., Oakey N.S., Hebert D., Platt T., 1986. Vertical nitrate fluxes in the oligotrophic ocean. Science, 234, 870-873.
- Licot M., 1985. Etude écohydrodynamique du front Liguro-Provençal au large de la Corse. Relation entre l'hydrodynamique, les paramètres physico-chimiques et la production primaire. Thèse de Doctorat, l'Université de Liège, 131 pp.
- Lohrenz S.E., Wiesenburg D.A., DePalma I.P., Johnson K.S., Gustafson D.E., 1988. Interrelationships among primary production, chlorophyll and environmental conditions in frontal regions of the Western Mediterranean Sea. Deep-Sea Res., 35, 793-810.
- Longhurst A.R., 1976. Interactions between zooplankton and phytoplankton profiles in the eastern tropical Pacific Ocean. Deep-Sea Res., 23, 729-754.
- Longhurst A.R., 1981. Significance of spatial variability. In A.R. Longhurst (Ed): Analysis of Marine Ecosystems. Academic Press London and New York, pp. 415-441.
- Longhurst A.R., 1989. Pelagic Ecology: Definition of pathways for material and energy flux. In M. Denis (Ed): Océanologie: actualité et perspective. Centre Océanologique de Marseille, pp. 263-288.
- Longhurst A.R., Sathyendranath S., Platt T., Caverhill C., 1995. An estimate of global primary production in the ocean from satellite radiometer data. J. Plankton Res., 17, 1245-1271.

- Longhurst A., Williams R., 1979. Materials for plankton modelling: vertical distribution of Atlantic zooplankton in summer. *J. Plankton Res.*, 1, 1-28.
- Lottka A.J., 1924. Contribution to the energetics of evolution. *Proc. Nat. Acad. Sci.*, 8, 147-150.
- Lutjeharms J.R.E., Walters M.N., Allanson B.R., 1985. Oceanic frontal systems and biological enhancement. In W.R. Sigafoos, P.R. Condy & R.M. Laws (Eds): *Antarctic Nutrient Cycles and Food Webs*. Springer-Verlag, Berlin Heidelberg New York, pp. 11-21.
- MacCarthy J.J., 1972. The uptake of urea by marine phytoplankton. *J. Phycol.*, 8, 216-222.
- MacCarthy J.J., 1981. The kinetics of nutrient utilisation. *Can. J. Fish. Aquat. Sci.*, Vol., 211-233.
- MacCarthy J.J., Carpenter E.J., 1983. Nitrogen cycling in near-surface waters of the open ocean. In: *Nitrogen in the marine environment*. Academic Press, pp. 487-512.
- MacIsaac J.J., Dugdale R.C., 1969. The kinetics of nitrate and ammonia uptake by natural populations of marine phytoplankton. *Deep-Sea Res.*, 16, 45-57.
- MacIsaac J.J., Dugdale R.C., 1972. Interactions of light and inorganic nitrogen in controlling nitrogen uptake in the sea. *Deep-Sea Res.*, 19, 209-232.
- Mann K.H.: 1969. The analysis of aquatic ecosystems. In R.B. Clark & R.J. Wooton (Eds): *Essays in Hydrobiology*. Univ. Exeter, U.K., pp. 1-14.
- Mann K.H., Lazier J.R.N., 1991. *Dynamics of Marine Ecosystems*. Blackwell Sci. Publ., 466 pp.
- Mantoura R.F.C., Llewellyn C.A., 1983. The rapid determination of algal chlorophyll and carotenoid pigments and their breakdown products in natural waters by reverse-phase high-performance liquid chromatography. *Anal. Chim. Acta*, 151, 297-314.
- Marchant H.J., Murphy E.J., 1994. In S.Z. El-Sayed S.Z. (Ed): *Southern Ocean Ecology: The Biomass Perspective*, pp. 267-285.
- Margalef R., Ballester A., 1967. Fitoplancton y producción primaria de la costa catalana, de junio de 1965 a junio de 1966. *Inv. Pesq.*, 31, 165 - 182.
- Margalef R., Castellvi J., 1967. Fitoplancton y producción primaria de la costa catalana, de julio de 1966 a julio de 1967. *Inv. Pesq.*, 31, 491-502.
- Marr J.W.S., 1962. The natural history and geography of the Antarctic krill *Euphausia superba* (Dana). *Disc. Rep.*, 32, 33-464.
- Marra I., Bidigare R.R., Dickey T.D., 1990. Nutrients and mixing, chlorophyll and phytoplankton growth. *Deep-Sea Res.*, 37, 127-143.
- Marshall S.M., Orr A.P., 1952. On the biology of *Calanus finmarchicus*. VII. Factors affecting eggs production. *J. Mar. Bio. Assoc. U.K.*, 30, 527-548.
- Marshall S.M., Orr A.P., 1972. The biology of a marine copepod. Springer Verlag, 195 pp.
- Martin J.H., Fitzwater S. E., Gordon R. M., Hunter C. N., Tanner S.J., 1993. Iron, primary production and carbon-nitrogen flux studies during the JGOFS North Atlantic Bloom Experiment. *Deep-Sea Res.*, 40, 115-
- Martin J.H., Gordon R.M., Fitzwater S.E., 1990. Iron in Antarctic waters. *Nature*, 345, 156-158.
- Martin J.H., Gordon R.M., Fitzwater S.E., 1991. The case for iron. *Limnol. Oceanogr.*, 36, 1793-1802.
- Mathieu T., 1992. Influence de la fonte de la glace marine sur la couche de mélange. Mémoire de Diplôme d'étude approfondie Européen en Modélisation de l'environnement marin, 82 pp.
- Mauclaire J. 1998. The biology of Calanoid copepods. Academic Press, San Diego, 710 pp.
- Maykut G.A., 1978. Energy exchange over young sea ice in the Central Arctic. *J. Geophys. Res.*, 83, 3646-3658.
- Maykut G.A., Perovich D.K., 1987. The role of shortwave radiation in the summer decay of a sea ice cover. *J. Geophys. Res.*, 92, 7032-7044.
- Maykut G.A., Untersteiner N., 1971. Some results from a time-dependent thermodynamic model of sea ice. *J. Geophys. Res.*, 76, 1550-1575
- Mayzaud P., 1973. Respiration and excretion of zooplankton. II. Studies of the metabolic characteristics of starved animals. *Mar. Biol.*, 21, 19-28.
- Mayzaud P., 1986. Digestive enzymes and their relation to nutrition. In E.D.S. Corner, S.C.M. O'Hara, (Eds): *The biological chemistry of marine copepods*. Clarendon, Oxford, pp 16-25.
- Mayzaud P., Dallot S., 1973. Métabolisme du zooplanton: respiration et excretion azotée. I étude des niveaux métaboliques de quelques espèces. *Mar. Biol.*, 19, 307-314.

- Mc Allister C.D., 1970. Zooplankton rations, phytoplankton mortality and the estimation of marine production. In J.H. Steele (Editor), *Marine Food Chains*. Oliver and Boyd, Edinburgh, 419-457.
- Mc Climans T.A., Lonseth L. 1985. Oscillations of frontal currents. *Cont. Shelf Res.*, 4, 699-707.
- Mellor G.L., Kantha L., 1989. An ice-ocean coupled model. *J. Geophys. Res.*, 94, 10937-10954.
- Melnikov I.A., 1998. Winter production of sea ice algae in the western Weddell Sea.. In J. Le Fevre and P. Trégouë (Eds): *Carbon Fluxes and Dynamic Processes in the Southern Ocean: Present and Past*. *J. Mar. Syst.*, 17, 195-205.
- Michaelis L., Menten M.L., 1913. Der Kinetic der Inverteinwirkung. *Biochem. Z.*, 49, 333-369.
- Millot C., 1979. Wind induced upwellings in the Gulf of Lions. *Oceanol. Acta*, 2, 261-274.
- Minas H.J., Minas M., Coste B., Gostan J., Nival P., Bonin M.C., 1988. Production de base et de recyclage: une revue de la problématique en Méditerranée Nord - Occidentale. *Oceanol. Acta*, 9 (Spec. Issue), 155-162.
- Mitchell B.G., Holm Hansen O., 1991a. Bio-optical properties of Antarctic Peninsula waters: differentiation from temperate ocean models. *Deep-Sea Res.*, 38, 1009-1028.
- Mitchell B.G., Holm Hansen O., 1991b. Observation and modeling of the Antarctic phytoplankton crop in relation to mixing depth. *Deep-Sea Res.*, 38, 981-1007.
- Mitchell J.G., Okubo A., Fuhrman J.A., 1985. Microzones surrounding the phytoplankton form the base for a stratified marine microbial ecosystem. *Nature*, 316, 58-59.
- Moloney C.L., Field J.G., 1991. The size-based dynamics of plankton food webs. I. A simulation model of carbon and nitrogen flows. *J. Plankton Res.*, 13, 1003-1038.
- Monzicoff A., Dallot S., Pizay M.D., 1992. Blue and yellow fluorescence of filtered seawater in a frontal zone (Ligurian Sea, northwest Mediterranean Sea). *Deep-Sea Res.*, 39, 1481-1498.
- Moore B., Bolin B., 1987. The oceans, carbon dioxide and global climate change. *Oceanus*, 28, 9-15.
- Morel A., André J.M., 1991. Pigment distribution and primary production in the Western Mediterranean as derived and modeled from coastal zone color scanner observations. *J. Geophys. Res.*, 96, 12685-12698.
- Mork M. 1981. Circulation phenomena and frontal dynamics of the Norwegian coastal current. *Phil. Trans. Soc. Lond.*, 302, 635-647.
- Mullin M.M., 1963. Some factors affecting the feeding of marine copepods of the genus *Calanus*. *Limnol. Oceanogr.*, 8, 239-250.
- Mullin M.M., 1991. Relative variability of reproduction and mortality in two pelagic copepod populations. *J. Plankton Res.*, 13, 1381-138.
- Mullin M.M., Brooks E.R., 1970. Growth and metabolism of two planktonic, marine copepods as influenced by temperature and type of food. In J.H. Steele (Ed): *Marine Food Chains*. Oliver and Boyd. Mullin M.M., Brooks E.R., 1970. Production of the planktonic copepod, *Calanus helgolandicus*. *Bull. Scripps Inst. Oceanogr.*, 17, 89-103.
- Mullin M.M., Stewart E.F., Fuglister F.J., 1975. Ingestion of planktonic grazers as a function of concentration of food. *Limnol. Oceanogr.*, 20, 259-262.
- Murray J.W., Downs J.N., Strom S., We, C.L., Jannasch H.W., 1989. Nutrient assimilation, export production and Th scavenging in eastern equatorial Pacific. *Deep-Sea Res.*, 36, 1471-1489.
- Murphy L.S., Haugen E.M., 1985. The distribution and abundance of phototrophic ultraplankton in the North Atlantic. *Limnol. Oceanogr.*, 30, 47-58.
- Myklestad S., 1977. Production of carbohydrates by marine planktonic diatoms. II. Influence of the N/P ratio in the growth medium on the assimilation ratio, growth rate, and production of cellular and extracellular carbohydrates by *Chaetoceros affinis* var *Willei* (Gran) Hustedt and *Skeletonema costatum* (Grev) Cleve. *J. Exp. Mar. Biol. Ecol.* 29, 161-179.
- Nagata T., Kirchman D.L., 1991. Release of dissolved free and combined amino acids by bacterivorous marine flagellates. *Limnol. Oceanogr.* 36, 433-443.
- Nelson D.M., DeMaster D.J., Dunbar R.B., Smith W.O.J., 1996. Cycling of organic carbon and biogenic silica in the Southern Ocean: estimates of water-column and sedimentary fluxes in the Ross Sea continental shelf *Journal of Geophysical Research*, 101, 18519-18532.

- Neumann G., Pierson W.J., 1966. Principles of Physical Oceanography. Prentice-Hall, Englewood Cliffs, New Jersey. 545pp
- Newell R.C., Linley E.A.S., 1984. Significance of microhétérotrophes in the decomposition of phytoplankton: estimates of carbon and nitrogen flow based on the biomass of plankton communities. Mar. Ecol. Prog. Ser., 16, 105-119.
- Nicholls A.G., 1933. On the biology of *Calanus finmarchicus*. III. Vertical distribution and diurnal migration in the Clyde Sea area. J. Mar. Biol. Assoc. U.K., 19, 139-164.
- Nicol J.A.C., 1960. The biology of marine animals. Pitman, London, 699pp.
- Nielsen T.G., Richardson K., 1989. Food chain structure of the North Sea plankton communities: seasonal variations of the role of the microbial loop. Mar. Ecol. Prog. Ser., 56, 75-87.
- Nihoul J.C.J., 1975. Passive dispersion models. In J.C.J. Nihoul (Ed): Modeling of Marine Systems. Elseviers Oceanography Series, Elsevier, New York, 10, 69-95.
- Nihoul J.C.J., 1980. Residual circulation, long waves and mesoscale eddies in the North Sea. Oceanol. Acta, 3, 309-316.
- Nihoul J.C.J., 1984. A three-dimensional general marine circulation model in a remote sensing perspective. Ann. Geoph., 2, 433-442.
- Nihoul J.C.J., 1985. Modélisation des processus physiques, chimiques et biologiques dans un estuaire: exemple de l'estuaire et de la plume de l'Escaut. Naturaliste Canadien (Rev. Ecol. Syst.), 112, 15-29.
- Nihoul J.C.J., 1986. Introduction à la mécanique des fluides géophysiques. Association Université de Liège-Université de Corse, 85 pp.
- Nihoul J.C.J. 1989. Les modèles mathématiques: Base indispensable à l'étude interdisciplinaire des systèmes marins. In M. Denis (Ed): Océanologie: actualité et perspective. Centre Océanologique de Marseille, pp. 187-211.
- Nihoul J.C.J., Pichot G., Hecq J.H., Dubois D., 1972. Exploitation des données et modelage du système marin. Modèle mathém. Poll. mer du Nord (CIPS): synthèse, 2, 9-23.
- Nihoul J.C.J., Ronday F., 1976. Hydrodynamic models of the North Sea. Mém. Soc. R. Sc. Liège, 10, 61-96.
- Nihoul J.C.J., Hecq J.H., 1984. Influence of the residual circulation on the physico-chemical characteristics of water masses and the dynamics of ecosystems in the Belgian coastal zone. Cont. Shelf Res., 3, 167-174.
- Nihoul J.C.J., Deleersnijder E., Djennidi S., 1989. Modelling the general circulation of shelf seas by 3D k-e models. Earth Sc. Rev., 26, 163-189.
- Nihoul J.C.J., Djennidi S., Hecq J.H., 1989. Modelling coastal/shelf systems with emphasis on long term trends. International Journal for Numerical Methods in Engineering, 27, 113-127.
- Nival P., Nival S., Leroy C., 1971. Essai d'utilisation de la méthode polarographique de dosage de l'oxygène pour l'estimation de la respiration d'*Acartia clausi*. Rapp. Comm.int Mer Médit., 20, 301-303.
- Nival P., Nival S., Palazzoli L., 1972. Données sur la respiration de différents organismes communs dans le plancton de Villefranche-sur-Mer. Mar. Biol., 17, 63-76.
- Nixon S.W., Kremer J.N., 1977. Narragansett Bay the development of a composite simulation model for a New England estuary. In: C. Hall, J. Day (Eds), Ecosystem Modeling in Theory and Practice: An Introduction with Case Histories. New York, Wiley, pp. 622-673.
- North Sea Task Force, 1993. North Sea Sub region 1. Assessment report. Norway, United Kingdom.
- Nuccio C., Innamorati M., Lazzara L., Mori G., Massi L., 1999. Spatial and temporal distribution of phytoplankton assemblages in the Ross Sea. In F. Faranda, L. Guglielmo & A. Ianora A. (Eds): Ross Sea Ecology, Springer, Berlin, Heidelberg, pp. 231-245.
- Omori M., 1969. Weight and chemical composition of some important oceanic zooplankton in the North Pacific Ocean. Mar. Biol., 3, 4-10.
- Oszoy E., Latif M.A., Unluata U., 1981. On the formation of Levantine intermediate water. Rapp. Comm. Int. Mer Médit., 27(6), 51-66.
- Owens N.J.P., Woodward E.M.S., Aiken J., Bellan I.E., Rees A.P., 1990. Primary production and nitrogen assimilation in the North Sea during July 1987. Neth. J. of Sea Res., 25, 143-154.

- Pace M.L., Glasser J.E., Pomeroy L.R., 1984. A simulation analysis of continental shelf food webs. Mar. Biol., 82, 47-63.
- Paffenhofer G.A., 1970. Cultivation of *Calanus helgolandicus* under controlled conditions. Helgolander Wiss. Meeres., 20, 346-359.
- Paffenhofer G.A., Knowles S.C., 1979. Ecological implications of fecal pellet size, production and consumption by copepods. J. Mar. Res., 37, 35-47.
- Pakhomov E.A., Perissinotto R., Fronemen P.W., 1999 Predation impact of carnivorous macrozooplankton and microneuston in the Atlantic sector of the Southern Ocean. J. Mar. Syst., 19, 47-64.
- Palmisano A.C., SooHoo, J.B. SooHoo S.L., Kottmeier S.T., Craft L.L., Sullivan, C.W., 1986. Photoadaptation in *Phaeocystis pouchetii* advected beneath annual sea ice in McMurdo Sound, Antarctica. J. Plankton Res., 8, 891-906.
- Parkinson C.L., Washington W.M., 1979. A large-scale numerical model of sea ice. J. Geophys. Res., 84, 311-337.
- Parson T.P., Kessler T.A., An ecosystem model for the assessment of plankton production in relation to the survival of young fish. J. Plankton Res., 37, 35-49.
- Parsons T.R., LeBrasseur R.J., Fulton J.D., 1967. Some observations on the dependence of zooplankton grazing on the cell size and concentration of phytoplankton blooms. J. Ocean. Soc. Jap., 23, 10-17.
- Parsons T. R., Takahashi M., 1973. Biological oceanographic processes. Pergamon, Press Toronto, 186pp.
- Pergent G.P., Mendez S., Pergent C.M., Pasqualini V., 1999. Preliminary data on the impact of fish farming facilities on *Posidonia oceanica* meadows in Mediterranean. Oceanol. Acta, 22, 95-107.
- Petipa T.S., 1967. On the efficiency of utilization of energy in pelagic ecosystems of the Black Sea. Fish Res. Bd. Can. Translation, 973.
- Petit B., Norro A. 2000. Seasonal evolution of sea ice and oceanic heat flux in the Weddell Sea. J. Mar. Syst., 27, 37-52.
- Philippe M., Hareng L., 1982. Surface temperature fronts in the Mediterranean Sea from infrared satellite imagery. In J.C.J. Nihoul (Ed): Hydrodynamics of semi-enclosed seas, Elsevier Oceanography Serie, Amsterdam, 13, 91-128.
- Pingree R.D., 1978. Cyclonic eddies and cross frontal mixing. J. Mar. Biol. Ass. U.K., 58, 955-963.
- Platt T., Gallegos C.L., Harrison W.G., 1980. Photoinhibition of photosynthesis in natural assemblages of marine phytoplankton. J. Mar. Res., 38, 687-701.
- Platt T., Mann K.H., Ulanowicz R.E. (Eds), 1981. Mathematical models in Biological Oceanography, UNESCO Monographs on Oceanographic Methodology, 7, 156 pp.
- Pollard R.T., 1977. Observations and theories of Langmuir circulations and their role in near surface mixing. In M. Angel (Ed), A Voyage of Discovery. Pergamon. 235-251 pp.
- Pomeroy L.R., Wiebe W.J., 1988. Energetics of microbial food webs. Hydrobiologia, 159, 7-18.
- Pond S., Packard G.L., 1983. Introductory dynamical oceanography. Pergamon Press, 2nd edition, 329 pp.
- Pondaven P., Fravalo C., Ruiz-Pino D., Trégouët P., Quéguiner B., Jeandel, C., 1998. Modelling the silica pump in the permanently open ocean zone of the Southern Ocean. In J. Le Fevre and P. Trégouët (Eds): Carbon Fluxes and Dynamic Processes in the Southern Ocean: Present and Past. J. Mar. Syst., 17-587-619.
- Postma H., Rommets J.W., 1984. Variations of particulate organic carbon in the central North Sea. Neth. J. Sea Res., 18, 31-50.
- Poulet S.A., 1983. Comparison between five coexisting species of marine copepods feeding on naturally occurring particulate matter. Limnol. Oceanogr., 23, 1126-1143.
- Poulet S.A., Ianora A., Miralto A., Meijer L., 1994. Do diatoms arrest embryonic development in copepods? Mar. Ecol. Prog. Ser., 111, 79-86.
- Prieur L., 1973. Interprétation d'une zone de forte turbidité sur la radiale Calvi-Nice. Rapp. Comm. Int. Mer Médit., 22, 59-61.

- Prieur L., 1981. Hétérogénéité spatio-temporelle dans le bassin Liguro-Provençal. Rapp. Comm. Int. Mer Médit., 27(6), 177-179.
- Prieur L., 1985. Systèmes frontaux en mer Ligure à partir des mesures multiparamétriques en continu. Courrier Mediprod, 2, 2-5.
- Prieur L., Tiberti M., 1985. Identification et échelles des processus physiques et biologiques responsables de l'hétérogénéité spatiale près du front de Mer Ligure. Rapp. Comm. int. Mer Médit., 29, 35-36.
- Prieur L., Legendre L. 1988. Oceanographic criteria for new phytoplankton production. In B.J. Rothschild (Ed): Toward a theory on biological-physical interactions in the world ocean. Kluwer Academic Publisher, pp. 72-112.
- Proctor L.M., Furman J.A., 1990. Viral mortality of marine bacteria and cyanobacteria. Nature, 342, 60-62.
- Rassoulzadegan F. 1990. Marine protozooplankton and food webs. Zool. Sci. 7(suppl.), 189-196.
- Rassoulzadegan F., 1993. Protozoan patterns in the Azam-Ammerman's bacteria-phytoplankton mutualism. In R. Guerrero & C. Pedrós-Alió (Eds): Trends in microbial ecology, Spanish Society for Microbiology, Barcelona, pp. 435-439.
- Raven J.A., Richardson K., 1986. Marine environments. In N.R. Baker, S.P. Long (Eds): Photosynthesis in contrasting environments. Elsevier, 337-396.
- Raymont J.E.G., 1963. Plankton and productivity in the oceans. Pergamon Press, Oxford, 1, 245 pp.
- Redfield A.C., Ketchum B.H., Richards F.A., 1963. The influence of organisms on the composition of sea water. In M.N. Hill (Ed): The Sea, ideas and observations on progress in the study of the seas, Interscience, New York, 26-77.
- Reid P.C., Lancelot C., Gieskes W.W.C., Hagmeier E., Weichert G., 1991. Phytoplankton of the North Sea and its dynamics: a review. Neth. J. Sea Res., 26, 295-331.
- Rey F., 1981. The development of the spring phytoplankton outburst at selected sites off the Norwegien coast. In: The Norwegian Coastal Current, University of Bergen, 2, 649-680.
- Ricard M., 1987. Atlas du phytoplancton marin: Diatomophycées. Editions du Centre National de la Recherche Scientifique, Paris, 297 pp.
- Richman S., Rogers J.N., 1969. The feeding of *Calanus helgolandicus* on synchronously growing populations of the marine diatom *Ditylum brightwelli*. Limnol. Oceanogr., 14, 701-709.
- Ricker W. E., 1975. Composition and interpretation of biological statistics of fish populations. Bull. Fish Res. Bd., 191, 1-382.
- Riley G.A., 1947. A theoretical analysis of the zooplankton population of Georges Bank. J. Mar. Res., 6, 104-113.
- Riley G.A., 1970. Particulate organic matter in seawater. Adv. Mar. Biol., 8, 11-18.
- Riley G.A., Stommel H., Bumpus D.F., 1949. Quantitative ecology of the plankton of the Western North Atlantic. Bull. Bingham. Oceanogr. Coll. 12, 1-169.
- Robertson J.R., 1983. Predation by estuarine zooplankton on tintinnid ciliates. Estuarine Coastal Shelf Sci., 16, 27-
- Robinson M.K., Bauer R.A., Schroeder E., 1979. Atlas of North Atlantic ocean monthly mean temperatures and mean salinities of the surface layer. US Naval Oceanographic Office Reference Publication, 18.
- Rodriguez-Prada R., 1973. Contribution à l'étude de la production primaire du bassin Liguro-Provençal. Thèse de 3^{me} cycle, Université Pierre et Marie Curie, Paris, 86pp.
- Rothschild B.J., Osborn T.R., 1988. Small-scale turbulence and planktonic contact rates. J. Plankton Res., 10, 465-474.
- Roy S., Harris R.P., Poulet S.A., 1989. Inefficient feeding by *Calanus helgolandicus* and *Temora longicornis* on *Coscinodiscus wailesii*: quantitative estimation using chlorophyll-type pigments and effects on dissolved free amino acids. Mar. Ecol. Prog. Ser. 52, 145-153.
- Ruiz J., Garcia C.M., Rodriguez J., 1996. Sedimentation loss of phytoplankton cells from the mixed layer: effects of turbulence levels. J. Plankt. Res., 18, 1727-1734.
- Russel F.S., 1931. The vertical distribution of marine macroplankton. X Further observations on diurnal changes. J. Mar. Biol. Assoc. U.K., 17, 767-784.

- Ryther J.H., Dunstan W., Tenore K., Huguenin J., 1972. Controlled eutrophication increasing food production from the sea by recycling human wastes. Biosci., 22, 144-152.
- Saggiomo V., Goffart A., Carrada G.C., Hecq J.H., 1994. Spatial patterns of phytoplanktonic pigments and primary production related with environmental parameters of a periantarctic ecosystem: The Strait of Magellan. J. Mar. Sys., 5, 119-142.
- Saint-Guily B., 1968. effet de la thermocline sur la diffusion d'une substance à partir d'une source. Vie et Milieu, 19, 1-8.
- Saiz E., Alcaraz M., 1991. Effects of small- scale turbulence on development time and growth of *Acartia grani* (Copepod: Calanoida). J. Plankton Res., 13, 873-883.
- Sakshaug E., Myklestad S., Andresen N.K., Hegseth E., Jorgensen L., 1981. Phytoplankton off the More coast in 1975-1979: distribution, species composition, chemical composition and conditions for growth. In: The Norwegian Coastal Current, University of Bergen, 2, 681-711.
- Sakshaug E., Slagstad D., 1991. Light and productivity of phytoplankton in polar marine ecosystem: a physiological view. In E. Sakshaug, C.C.E. Hopkins & N.A. Ørntsland (Eds): Proceedings of the Pro Mare Symposium on Polar Marine Ecology, Trondheim, 12-16 may 1990. Polar Res., 10, 69-85.
- Santer B., 1996. Nutritional suitability of the dinoflagellate *Ceratium furcoides* for four copepod species. J. Plankton. Res., 18, 323 - 333.
- Sarmiento J.L., Fasham M.J.R., Siegenthaler U., Najjar R., Toggweiler J.R., 1989. Models of chemical cycling in the ocean: Progress Report II. Ocean Tracer Laboratory Technical Report n°6, Princeton University.
- Satoh H., Watanabe K., Khanda H., Takahashi E., 1986. Seasonal changes of chlorophyll *a* standing stocks and oceanographic conditions under fast ice near Syowa Station, Antarctica. Antarctic record, 30, 19-32.
- Savage M.L., Stearns C., 1985. Climate in the vicinity of Ross Island, Antarctica. Antarctic Journal of the United States, 20, 1-9.
- Savenkoff C., Prieur L., Reys J.P., Lefèvre D., Dallot S., Denis M., 1993. Deep microbial communities evidenced in the Liguro-Provençal front by their ETS activity. Deep-Sea Res., 40, 709-725.
- Scholander P.F., Flagg W., Walters V., Irving L., 1953. Climatic adaptation in arctic and tropical poikilotherms. Physiol. Zool., 26, 67-92.
- Sedwick P.N., DiTullio G.R., 1997. Regulation of algal blooms in Antarctic shelf waters by the release of iron from melting sea ice. Geophys. Res. Lett., 24, 2515-2518.
- Semtner A.J. Jr , 1976. A model for the thermodynamic growth of sea ice in numerical investigations of climate. J. Geophys. Res., 6, 379-389.
- Shine K.P., Henderson-Sellers A., 1985. The sensitivity of a thermodynamic sea ice model to changes in surface albedo parameterization. J. Geophys. Res., 90, 2243-2250.
- Sieburth J. McN., 1960. Acrylic acid, an "antibiotic" principle in *Phaeocystis pouchetii* in Antarctic waters. Science, 132, 667-676.
- Sieburth J. McN., Smetacek V., Lenz J., 1978. Pelagic ecosystem structure: Heterotrophic compartments of the plankton and their relationship to plankton size fractions. Limnol. Oceanogr., 23, 1256-1263.
- Signorini S., McClain C., Christian J., Wong C.S., 2000 One-dimensional coupled ecosystem-carbon flux model for the simulation of biogeochemical parameters at ocean weather station P. Nasa Center for aerospace information. Nasa/TP-2000-209892, 36 pp.
- Skliris N., Elkalay K., Frangoulis C., Goffart A., Hecq J.H., 1999. A model approach of the plankton ecosystem of the north-western Corsican coastal area. Actes des 3èmes rencontres d'hydrodynamique marine. Univ. Ben M'Sik Press, Casablanca. In press.
- Skliris N., Elkalay K., Goffart A., Frangoulis C., Hecq J.H., 2001. One-dimensional modelling of the plankton ecosystem of the north-western Corsican coastal area in relation to meteorological constraints. J. Mar. Syst., 27, 337-362.
- Skliris N., Goffart A., Hecq J.H., Djenidi S., 2001. Shelf-slope exchanges associated with a steep submarine canyon off Calvi (Corsica, NW Mediterranean Sea): a modelling approach. In press in J. Geophys. Res.

- Small F.L., Fowler S.W., Unlu M.Y., 1979. Sinking rates of natural copepod fecal pellets. Mar. Biol., 51, 233-241.
- Smayda T.J., 1970. The suspension and sinking of phytoplankton in the sea. Oceanogr. Mar. Biol. Ann. Rev., 8, 353-414.
- Smayda T.J. 1971. Normal and accelerated sinking of phytoplankton in the Sea. Marine Geol., 11, 105-122
- Smetacek V., 1981. The annual cycle of protozooplankton in the Kiel Bight. Mar. Biol., 63, 1-8
- Smetacek V., Scharek R., Nöthig E.M., 1990. Seasonal and regional variation in the pelagic, and its relationship to the life history cycle of krill. In K.R. Kerry, G. Hempel (Eds): Antarctic Ecosystems: Ecological Change and Conservation. Springer Verlag Berlin, pp 103-114.
- Smith W.O., Codispodi L.A., Nelson D., Manley T., Buskey E.J., Niebauer H.J., Cota G.F., 1991. Importance of *Phaeocystis* blooms in the high-latitude ocean carbon cycle. Nature, 352, 514-516.
- Smith W.O., Nelson D.M., 1985. Phytoplankton bloom produced by a receding ice edge in the Ross Sea: spatial coherence with the density field. Science, 227, 163-166.
- Smith W.O., Gordon L.I., 1997. Hyperproductivity of the Ross Sea (Antarctica) polynya during austral spring. Geophys. Res. Lett., 24, 223-236.
- Sournia A., 1969. Cycle annuel du phytoplancton et de la production primaire dans les mers tropicales. Mar. Biol., 3, 287-303.
- Sournia A., Brylinski J.M., Dallot S., Le Corre P., Leveau M., Prieur L., Froget C., 1990. Fronts hydrologiques au large des côtes françaises: Les sites-ateliers du programme Frontal. Oceanol. Acta, 13, 413-438.
- Stavn R.H., 1971. The horizontal, vertical distribution hypothesis: Langmuir circulation and *Daphnia* distributions Limnol. Oceanogr., 16, 658-664.
- Steele J.H., 1958. Plant production in the northern North Sea. Marine Res. (Scot Home Dept), 7, 1-36.
- Steele J.H., 1974. The structure of marine ecosystem. Blackwell Scientific Publications, 128 pp.
- Steele J.H., Henderson E.W., 1992. The role of predation in plankton models. J. Plankton Res., 14, 157-172.
- Steeman Nielsen E., 1975. Marine photosynthesis with special emphasis on the ecological aspects. Elsevier Oceanography Series, Amsterdam:
- Steidinger K.A., Walker L.M., 1984. Introduction. In K.A. Steidinger and L.M. Walker (Eds): Marine Plankton Life Cycle Strategies, CRC Press Boca Raton, Fla, 1-4
- Stoecker D.K., Capuzzo J., 1990. Predation on protozoa: its importance to zooplankton. J. Plankton Res., 12, 891-898.
- Stoecker D.K., Sanders N.K., 1985. Differential grazing by *Acartia tonsa* on a dinoflagellate and a tintinniid. J. Plankton Res., 7, 85-100.
- Strickland J.D.H., 1960. Measuring the production of marine phytoplankton. Fish. Res. Bd. Can. Bull., 122, 1-172.
- Sturman A.P., Anderson M.R., 1986. On the sea-ice regime of the Ross Sea, Antarctica. J. Glaciology, 32, 54-59.
- Sullivan C.W., McClain C.R., Comiso J.C., Smith, W.O., 1988. Phytoplankton standing crops within an Antarctic ice edge assessed by satellite remote sensing. J. Geophys. Res., 93, 12487-12498.
- Sugimura Y., Suzuki Y., 1988. A high-temperature catalytic oxidation method of non-volatile dissolved organic carbon in seawater by direct injection of liquid samples. Mar. Chem., 14, 105-131.
- Suijlen J.M., Van LeussenW., 1990. Recent developments in large-scale tracing by fluorescent tracers. In W. Michaelis (Ed): Estuarine Water Quality Management. Coastal and Estuarine Studies, Springer Verlag, Berlin, 36, pp181-188.
- Sverdrup H.A., 1953. On conditions for the vernal blooming of phytoplankton. J. cons. int. perm. explor. Mer, 18, 287-295.
- Swanberg N.R., 1985. The trophic role of colonial radiolaria in oligotrophic oceanic environments. Limnol. Oceanogr., 28, 655-666.

- Tande K.S., Båmstedt U., 1987. On the trophic fate of *Phaeocystis pouchetii*. 1. Copepod feeding rates on solitary cells and colonies of *P. pouchetii*. Sarsia, 72, 313-320.
- Tanimura A., Hoshiai T., Fukuchi M., 1996. The life cycle strategy of the ice-associated copepod, *Paralabidocera antarctica* (Calanoida, Copepoda) at Syowa Station, Antarctica. Antarctic Science, 8, 257-266.
- Taylor G.T., 1982. The role of pelagic heterotrophic protozoa in nutrient cycling: a review. Ann. Inst. Océanogr., Paris, 58(S), 227-241.
- Taylor A.H., Joint I., 1990. A steady-state analysis of the 'microbial loop' in stratified systems. Mar. Ecol. Prog. Ser., 59, 1-17.
- Taylor A.H., Watson A.J., Ainsworth M., Robertson J.E., Turner D.R., 1991. A modelling investigation of the role of phytoplankton in the balance of carbon at the surface of the North Atlantic. Global Biogeochem. Cycles, 5, 151-171.
- Taylor A.H., Harbour D.S., Harris R.P., Burkhill P.H., Edwards E.S., 1993. Seasonal succession in the pelagic ecosystem of the North Atlantic, the utilization of nitrogen. J. Plankton Res., 15, 875-891.
- Taylor A.H., Geider R.J., Gilbert F.J.H., 1997. Seasonal and latitudinal dependencies of phytoplankton carbon to chlorophyll *a* ratios: results of a modelling study. Mar. Ecol. Prog. Ser., 152, 51-66.
- Tchernia P., 1980. Descriptive Physical Oceanography. Pergamon Press, Oxford, 253pp.
- Tegen I., Fung I., 1994. Modeling of mineral dust in the atmosphère: Sources, transport and optical thickness. J. Geophys. Res., 99, 22897-22914.
- Tett P., 1981. Modelling phytoplankton production at shelf-sea fronts. Phil.Trans. r .Soc. Lond. A., 302, 605-615.
- Tett P., Edwards A., Jones K., 1986. A model for the growth of shelf-sea phytoplankton in summer. Estuar. Coast. Shelf Sci., 23, 641-472.
- Therriault J.C., Lauwrence D.J., Platt T., 1978. Spatial variability of phytoplankton turnover in relation to physical processes in a coastal environment. Limnol. Oceanogr., 23, 900-911.
- Thingstad T.F., Rassoulzadegan F., 1995. Nutrient limitations, microbial food webs, and 'biological C-pumps': suggested interactions in a P-limited Mediterranean. Mar. Ecol. Prog. Ser., 117, 299-306.
- Thingstad T.F., Skjoldal E.F., Bohne R.A., 1993. Phosphorus cycling and algal-bacterial competition in Sandefjord, western Norway. Mar. Ecol. Prog. Ser., 99, 239-259.
- Thomas J.P., 1971. Release of dissolved organic matter from natural populations of marine phytoplankton. Mar. Biol., 11, 311-323.
- Thorson G., 1979. Havbundensdyreliv. Infraunaen, den jaevne havbunds dyresamfund. In A Norrevang , J. Lundo: Danmarks natur, Politikens forlag, Copenhagen, 3, pp 82-157.
- Thurman H.V., Weber H.H., 1984. Marine Biology, Charles E. Merrill, Columbus, Ohio, 382 pp
- Toggweiler J.R., 1989. Is the downward dissolved organic matter (DOM) flux important in carbon transport? In W.H. erger, V.S. Smetacek, G. Wefer (Eds): Productivity in the Oceans: Present and Past, Wiley, 65-83.
- Touratier F., 1996. Modélisation Ecologique en Région Septentrionale de la Mer du Nord : Application à la campagne de Mesures du Fladen Ground 1976 (Flex'76). Thèse de Doctorat, Université de Liège, 302 pp.
- Travers A., Travers M., 1973. Données sur quelques facteurs de l'écologie du plancton dans la région de Marseille. 3. La lumière. Téthys, 5, 7-30.
- Tréguer P., Jacques G., 1992. Dynamics of nutrients and phytoplankton, and fluxes of carbon, nitrogen and silicon in the Antarctic Ocean. Polar Biol., 12, 149-162.
- Tréguer P., Nelson D.M., Van Bennekom A.J., DeMaster D.J., Leynaert A., Quéguiner B., 1995. The silica balance in the World Oceans: a reestimate. Science, 268, 375-379.
- Turner J.T., 1977. Sinking rates of fecal pellets from the marine copepod *Pontella meadii*. Mar. Biol., 40, 249-259.
- Turrel W.R., Henderson E.W., Slesser G., Payne R., Adams R.D., 1992. Seasonal changes in the circulation of the northern North Sea. Cont. Shelf Res., 12(2-3), 257-286.

- Tusseau M.H., Lancelot C., Martin J.M., Tassin B., 1997. 1-D coupled physical-biological model of the northwestern Mediterranean Sea. Deep-Sea Res., 44, 851-880.
- Valiela I., 1995. Marine Ecological Processes, Springer Verlag, New York. 686 pp.
- Varela R.A., Crusado A., Tintoré J., 1994. A simulation analysis of various biological and physical factors influencing the deep-chlorophyll maximum structure in oligotrophic areas. J. Mar. Syst., 5, 143-157.
- Varela R. A., Crusado A., Tintoré J., Ladona E.G., 1992. Modelling the deep-chlorophyll maximum: A coupled physical-biological approach. J. Mar. Res., 50, 441-463.
- Veeschkens C., Hecq J.H., 1996. Modelling Approach of the Antarctic ecosystem. In Royal Academy of Belgium (Ed.): Progress in Belgian Oceanographic Research. Proceedings of the annual workshop on Belgian Oceanographic Research, 175-178.
- Veeschkens C., Belkhiria S., Goffart A., Hecq J.H., 1994. Distribution of plankton parameters in the North Eastern North Sea in relation to vertical physical structures. Bull. Soc. R. Sc. Liège, 63, 65-88.
- Verity P.G., 1985. Grazing, respiration, excretion and growth rates of tintinnids. Limnol. Oceanogr., 30, 1268-1282.
- Verity P.G., 1986. Growth rates of natural tintinnid populations in Narragansett Bay, Mar. Ecol. Prog. Ser., 29, 117-131.
- Verity P.G., Smayda T.J., 1989. Nutritional value of *Phaeocystis poucheti* (Primnesiophyceae) and other phytoplankton for *Acartia* spp. (Copepoda): ingestion, egg production and growth of nauplii. Mar. Biol., 100, 161-171.
- Vernet M., Lorenzen C.J., 1987. The relative abundance of pheophorbide α and pheophytin α in temperate marine waters. Limnol. Oceanogr., 32, 352-358.
- Veth C., 1991. The evolution of the upper water layer in the marginal ice zone, austral spring 1988, Scotia-Weddell Sea. J. Mar. Syst., 2, 451-464.
- Vinogradov M.E., 1970. Vertical distribution of the oceanic zooplankton. Israel program for Scientific translations, Jerusalem, 339 pp.
- Virtue P., Nichols P.D., Nicol S., McMinn A., Sikes E.L., 1993. The lipid composition of *Euphausia superba* Dana in relation to the nutritional value of *Phaeocystis pouchetii* (Hariot) Lagerheim. Antarctic Sciences 5, 169-177.
- Volterra V., 1926. Fluctuation in the abundance of a species considered mathematically. Nature, 188, 558-560.
- Veronina N.M., 1998. Comparative abundance and distribution of major filter-feeders in the Antarctic pelagic zone. In J. Le Fevre and P. Tréguer (Eds): Carbon Fluxes and Dynamic Processes in the Southern Ocean: Present and Past. J. Mar. Syst., 17, 375-390.
- Walsh J.J., 1975. A spatial simulation model of the Peru upwelling ecosystem. Deep-Sea Res., 22, 201-236.
- Walsh J.J., 1988. On the nature of continental shelves. Academic Press San Diego, 520 pp.
- Walsh J.J., Dugdale R.E., 1971. A simulation model of the nitrogen in the Peruvian upwelling system. Invest. Pesq., 35, 309-330.
- Watanabe K., Nakajima Y., 1982. Vertical distribution of chlorophyll α along 45°E in the Southern Ocean, 1981. Memoirs National Institute of Polar Research, Special Issue, 23, 73-86.
- Weller R.A., Dean J.P., Marra J., Price J.F., Francis E.A., Bordman D.C., 1985. Three-dimensional flow in the upper ocean. Science, 227, 1552-1556.
- Welschmeyer N.A., Lorenzen C.J., 1985. Chlorophyll budgets: zooplankton grazing and phytoplankton growth in a temperate fjord and the Central Pacific Gyres. Limnol. Oceanogr., 30, 1-21.
- Wheeler P.A., Kirschman D.L., 1986. Utilization of inorganic and organic nitrogen by bacteria in marine systems. Limnol. Oceanogr., 31, 998-1009.
- Williams P.J. leB., 1975. Biological and chemical aspects of dissolved organic material in seawater. In J.P. Ryley, G. Skirrow (Eds): Chemical Oceanography. Academic Press, New York, pp301-363.

- Williams P.J. Le B., 1990. The importance of losses during microbial growth: commentary on the physiology, measurement and ecology of the release of dissolved organic material. *Mar. Microb. Food Webs*, 4, 175-206.
- Williams R., 1984. An overview of secondary production in pelagic ecosystems. In M.J.R. Fasham (Ed): Flows of energy and materials in marine ecosystems. Theory and Practice. Nato Conf. Series, Plenum, 13, 361-405.
- Williams R., Claustre H., 1991. Photosynthetic pigments as biomarkers of phytoplankton populations and processes involved in the transformation of particulate organic matter at the Biotrans site (47°N, 20°W). *Deep-Sea Res.* 38, 347-355.
- Wilmotte A., Demonceau C., Goffart A., Hecq J.H., Demoulin V., Crossley A.C., 2001. Molecular and pigment studies of the picophytoplankton in a region of the Southern Ocean (47 to 54° S, 141 to 144 °E) in March 1998. Submitted to Deep-Sea Res.
- Wróblewski J., 1977. A model of phytoplankton plume formation during variable Oregon upwelling. *J. Mar. Res.*, 35, 357-394.
- Wróblewski J.S., Sarmiento J.L., Flierl G.R., 1988. An ocean basin scale model of plankton dynamics in the North Atlantic. 1. Solutions for the climatological oceanographic conditions in May. *Global Biogeochem. Cycles.*, 2, 199-218.
- Yentsch C.S., Lee R.W., 1966. A study of photosynthetic light reactions, and a new interpretation of sun and shade phytoplankton. *J. Mar. Res.*, 24, 319-337.
- Zwally H.T., Parkinson C.L., Comiso J.C., 1983. Variability of Antarctic sea ice and changes in carbon dioxide. *Science*, 220, 1005-1012.