

CARBONIFEROUS AND PERMIAN ALGAL MICROFLORA, TARIM BASIN (CHINA)

Bernard MAMET¹ & Zili ZHU²

1. *Université Libre de Bruxelles, Department of Earth and Environmental Sciences, 50 av. F.P. Roosevelt, CP 160/02, B-1050 Brussels*

2. *Nanjing Institute of Geology and Palaeontology, Academia Sinica, Chi-Ming-Ssu, Nanjing, China 210008*

(6 figures)

ABSTRACT. Thirty eight algal genera have been observed in Tournaisian to Kungurian carbonates of the Tarim Basin (China). Floral assemblages permit identification of ten stratigraphic levels. Most of the widespread and common forms have a “cosmopolitan” distribution, although *Calcifolium* is recognized for the first time in the eastern part of the Tethys.

Key-words: Tarim basin, Carboniferous, Permian, algae, China.

RESUME. Trente-huit genres d'algues calcaires ont été reconnus dans le Paléozoïque Supérieur du Bassin de Tarim (Chine). Les assemblages de flores permettent l'identification de dix niveaux stratigraphiques locaux. La plupart des taxa ont une distribution “cosmopolite”, à l'exception de *Calcifolium* qui est observé pour la première fois dans la partie orientale de la Téthys.

Mots-clés: Bassin de Tarim, Carbonifère, Permien, algues, Chine.

1. Introduction

This paper reports that extensive Carboniferous and Permian marine deposits in the Tarim Basin contain abundant calcareous algae. Previously, Mu (1985) reported in the Tianshan belt an Asselian to Artinskian flora with 13 genera ascribed to the Cyanophyta and the Chlorophyta. But little is known on the Carboniferous algae in this area, although they are present in many horizons. The material described here was collected by one of the authors, Zili Zhu, who joined the Upper Paleozoic research team organized by the Nanjing Institute of Geology and Palaeontology, Academia Sinica in 1992-1994. Recent oil exploration has made the general geology and stratigraphy of the basin better understood (Wang et al., 1992, Zhou & Chen, 1992, Zhai & Sun, 1994, Zhu, 1995, Liao & Rodriguez, 1999).

2. General geology

The Tarim Basin (Fig. 1) is a Proterozoic plate separated from other terranes by the Tianshan belt to the north and the Kunlun belt to the south. From the late Precambrian to the Ordovician, the northeastern part of the plate formed a southwest-northeast trending platform-basin system. This pattern gradually disappeared during the Silurian, Devonian, Carboniferous time and in the Permian, it finally resulted in the welding of the Tarim Basin

and the Kazakhstan-Dzungarian Block (southern part of the Asia plate) (Jia et al., 1992b). Consequently, the depositional center shifted to the western end of the basin in Carboniferous and Early Permian time (Wang, 1986). The marine history of the region ended in the Middle Permian (Early Guadalupian) (Zhou & Chen, 1992).

Major transgressions occur at three levels. The first is short and relatively local. In the Tournaisian, the sea invades the middle part of the basin to form the crinoidal limestone member of the Bachu Formation. This transgression comes from the north, and the southwestern part of the basin remains littoral. Open marine environments in the southwestern part of the basin are restricted to the upper part of the Kelitake Formation.

The second transgression is still local and begins in the early Viséan when the sea invades the northwestern and southeastern margins of the basin. The transgression comes from the west and lasts up to the Serpukhovian. It ended with a major regression at the base of the Kalawuyi Formation.

The third transgression begins in the Bashkirian and open marine conditions prevail in the western and middle part of the basin in Moscovian time. In the Early Permian, open marine facies cover most of the western and middle part of the basin. Those marine conditions end in Early Guadalupian time. Groves and Brenckle (1997) suggest an additional important Sakmarian hiatus, that is not corroborated here by the fusulinid succession.




Figure 1. Carboniferous and Early Permian paleogeography, Tarim Basin. In Carboniferous and Early Permian time the eastern part of the basin was eroded, while the central part was occupied by alternations of supersaline lagoons and open marine ramps. The western part of the basin, in particular the southwestern and northwestern margins, was characterized by open marine environments. Subsidence was active and an extensive and thick marine sequence of nearly 2500m was deposited.

3. General stratigraphy

The thick marine sequences of the western part of the basin contain diverse fossil assemblages that have been recorded by Zhao et al. (1984), by the Institute of Geology, Xinjiang Geological Bureau & Institute of Geology, CAGS (1987), by Zhang & Gu (1991) and Liao et al. (1992). Recent oil exploration encouraged further studies of the basin and a fine framework of stratigraphy was established (Fig. 2). The southwestern margin of the basin, where an extensive marine succession from the basal Carboniferous to the Middle Permian is exposed, was extensively studied and used as a standard for regional correlations. Zhao et al. (1984) divided the succession into seven formations, in ascending order Kelitake, Heshilafu, Kalawuyi, Azigan, Tahaqi, Keziliqiman, and Qipan (Fig. 3).

The Kelitake Formation conformably overlies the upper Devonian terrestrial Qizilafu Formation and is divided in two parts. The lower part comprises dolomitic limestones and argillaceous limestones with *Cyrtospirifer* and *Tenticospirifer*. The upper part consists of oolitic limestones interbedded with lime mudstones. The Heshilafu Formation is composed of shales and sandstones with limestone intercalations. The base of the Viséan is marked by a coarse conglomerate. Brachiopods, corals and foraminifera are present. The Kalawuyi Formation has a somewhat similar rock composition but contains more limestone levels. No extensive conglomerates have been found except one layer of small pebbles, the base of which marks the Bashkirian widespread regression. One foraminiferal zone is missing at the base of the formation which may suggest an unconformity, although no obvious hiatus is recognized in the field. Fusulines, brachiopods

	Stage	Southwestern Margin (Zhao et al., 1984)	Northwestern Margin (IGXGB & IG,CAGS, 1987)	Middle part of the Basin (Jia et al., 1992)
P	Guadalupian			
	Kungurian	Qipan Fm.		
	Artinskian	Keziliqiman Fm.	Sarazheyi Fm. Kunkelaqi Fm. Kake Fm.	Xiaohaizi Fm.
	Sakmarian- Asselian	Tahaqi Fm.	Zaerjake Fm.	
Gzhelian- Kasimovian	? ? ?			
C	Moscovian	Azigan Fm.	Bijingtawu Fm.	Kalashayi Fm.
	Bashkirian	Kalawuyi Fm.		
	Serpukhovian	Heshilafu Fm.	Kongtaikinggou Fm.	
	Visean		Mongdaleke Fm.	
	Tournaisian	Kelitake Fm.		Bachu Fm.
D	Famennian	Qizilafu Fm.		

Figure 2. Carboniferous and Early Permian stratigraphy in Tarim Basin.

and corals are common. The overlying Azigan Formation includes light colored, massive limestones. Fusulines are abundant in the lower part but rare higher up. The Tahaqi Formation has essentially the same aspect and fusulines dominate the fauna. The Keziliqiman Formation yields thin limestone, dolomitic limestone and sandstone layers. Brachiopods, fusulines and fenestellid bryzoans are the main fossils. The Qipan Formation is composed of alternations of thin limestone, shales and sandstone with abundant bryzoans, bivalves and brachiopods.

4. Microfacies and foraminifers

The following is a concise description of the microfacies and the associated foraminifers ("endothyrids" and "fusulinids"). Foraminiferal zones are those of Mamet (1974).

4.1. Kelitake Formation (also transliterated Kelitag Formation)

Thin-bedded, black, shaly, dolomitic limestones. Black, thin-bedded argillaceous mudstones. Light gray, pelloid grainstones. Medium- to thick-bedded bioclastic-oolitic packstones. Some evaporites. Mainly restricted facies with

few foraminifers and abundant calcispheres at the base. Marine ramp at the top.

Tournaisian (probably Zone 7?): *Brunsia*, *Septaglomospiranella*, *Tournayella*, *Tournayellina*.
Early Visean (Zone 11) : *Brunsia*, *Dainella*, *Earlandia*, *Globoendothyra*, *Latiendothyra*, *Priscella*, *Pseudoammodiscus*.

4.2. Heshilafu Formation

Fine-grained conglomerates and sandstones, overlain by black, thin-bedded bioclastic grainstones. Dark shales with minor black, thin-bedded, sandy grainstones. Black, thick-bedded, cherty grainstones/packstones, with abundant corals and brachiopods (Liao & Rodriguez, 1999). First levels of oncolites.

Middle Visean (Zone 13): *Archaeodiscus*, *Brunsia*, *Endothyra*, *Endothyranopsis*, *Eostaffella*, *Mediocris*, *Planoarchaeodiscus*, *Priscella*, *Pseudoammodiscus*.

Latest Visean (or slightly younger?)(Zone 16) : *Archaeodiscus*, *Climacammina*, *Endothyra*, *Endothyranopsis*, *Eostaffella*, *Neoarchaeodiscus*, *Palaeotextularia*, *Planospirodiscus*, *Pseudotaxis*, *Tetrataxis*.

Serpukhovian (Zone 18) : *Asteroarchaeodiscus*, *Biseriella*, *Climacammina*, *Earlandia*, *Endothyra*, *Endothyranopsis*,

Permian	Kungurian	Qipan Fm. —300m	Langella	
	Artinskian	Keziliqiman Fm. —450m	Eoparafusulina	
	Asselian-Sakmarian	Tahaqi Fm. —250m	Sphaeroschwagerina-Schwagerina-Pseudoschwagerina	
Carboniferous	Kasimovian-Ghzelian		Triticites	
		?	?	
	Moscovian	26	Azigan Fm. —250m	Fusulina Fusulinella
		25		Syzrania-Pseudopalaeospiroplectammina
		24		
	Bashkirian	22	Kalawuyi Fm. —550m	Ozawainella-Profusulinella
		21		Globivalvulina
	Serpukhovian	19		Bianella-Eostaffella-Loeblichia
		18		
		17	Heshilafu Fm. —450m	Neoarchaediscus-Planospirodiscus
Vissean	16		Archaediscus-Eostaffella	
	10		Dainella-Globoendothyra	
Tournaisian	9			
	8	Kelitake Fm. —300m	Tourmayella-Tourmayellina	

Figure 3. Detailed stratigraphy of the southwestern Tarim Basin, microfacies and foraminifers. The zones of the seven formations are based on small foraminifers and fusulines. The lower and middle part of the Carboniferous yield rather precise zones that can be correlated with other basins of the Tethys (Mamet, 1974; Pinard & Mamet, 1998). The Permian has no formal foraminifer zonation and only stages or series can be recognized.

Eostaffella, Globoendothyra, Loeblichia, Neoarchaediscus, Omphalotis, Palaeotextularia, Planospirodiscus, Pseudoammo-discus, Pseudoendothyra, Pseudotaxis, Tetrataxis.

4.3. Kalawuyi Formation

At the base, fine- to coarse-grained sandstones interbedded with thin foraminifer grainstones. Then, massive alternations of sandstones and limestones. Few algal bafflestones. At the top, massive fusuline grainstones. Mainly detrital, grading to open marine ramp.

Late Bashkirian (Zone 21 and 22: no evidence of zone 20 which is probably missing): Bradyina, Climacammina, Endothyra, Endothyranella, Eoschubertella, Globivalvulina, Ozawainella, Palaeotextularia, Profusulinella, Pseudoendothyra, Pseudostaffella, Tetrataxis, Verella.

Early Moscovian (Zone 23): Aljutovella, Biseriella, Bradyina, Bradyinelloides?, Climacammina, Deckerella, Eofusulina, Globivalvulina, Ozawainella, Palaeotextularia, Profusulinella, Pseudobradyna, Pseudopalaeospiroplectammina, Pseudostaffella, Syzrania.

4.4. Azigan Formation

Light-colored, homogeneous, massive packstones. Then interfingerings of dark/white limestones. Massive grainstones with boundstones at the top.

Middle Moscovian (Zone 24 and 25) Beedeina, Bradyina, Bradyinelloides, Climacammina, Endothyranella, Fusulinella, Globivalvulina, Hemidiscus, Pseudopalaeospiroplectammina, Pseudobradyna, Tetrataxis.

Late Moscovian (Zone 26): Bradyina, Bradyinelloides, Climacammina, Fusulina, Fusulinella, Tetrataxis.

4.5. Tahaqi Formation

Thin-bedded varicolored grainstones and dolomites, massive oncolithic grainstones.

Kasimovian-Ghzelian (no typical Kasimovian fusulines identified) : Bradyina, Climacammina, Quasifusulina, Carbonoschwagerina, Triticites.

Asselian-Sakmarian: Climacammina, Nodosinelloides, Protonodosaria, Pseudoschwagerina, Schwagerina, Sphaeroschwagerina, Zellia (at top).

4.6. Keziliqiman Formation

Light-colored, massive, oolitic, oncolithic grainstones, thin-bedded evaporites, dolomites, black, thin-bedded wackestones.

Artinskian: Eoparafusulina, Geinitzina, Nankinella, Nodosinelloides, Protonodosaria, Staffella, Syzrania.

4.7. Qipan Formation

Thin-bedded wackestones, high-energy grainstones with oncolithes interbedded with minor shales, sandstones and bindstones.

Kungurian (or slightly younger): Climacammina, Geinitzina, Globivalvulina, Langella, Nodosinelloides, Protonodosaria, Syzrania.

5. Algal microflora (Fig. 4)

Algae are too facies sensitive and too long-ranging to permit the establishment of a precise stratigraphic zonation. Carboniferous and Permian stratigraphy must rely on ammonoids, corals, conodonts or foraminifers. However, on a local basis, the Tarim microflora allows recognition of 10 assemblages that are locally useful. When the fauna is scarce or absent, the algal flora proves to be an important addition to the stratigraphy.

5.1. The calcisphere flora (Fig. 6A)

Abundant and diversified calcispheres (Fig. 6A) characterize the Tournaisian part of the Kelitake Formation. Cal-




Figure 4. Detailed stratigraphic distribution of calcareous algae and incertae sedis, with numbered algal floras, in the Tarim Basin, subdivided according to stages and forams zones.

calcspheres are algal spore kysts and proliferate in lagoons. They are cosmopolitan in the Paleozoic world and range from Devonian to Carboniferous (Mamet, 1991, 1998).

5.2. The primitive *Koninckopora* flora (Fig. 5G)

First representatives of the dasycladale *Koninckopora* are small with a poorly calcified cortex (Nakai & Kato, 1981). Although Wright (1981) has challenged the validity of the concept, it appears however quite constant and synchronous on a world-wide basis. It may be used to underline the early Visean.

5.3. The advanced *Koninckopora* flora

Massive, well-calcified, double-layered *Koninckopora* are observed in all Middle-Late Visean shallow-water open-marine grainstones. They form an essential part of the "oolite factory". In Europe the taxon peters out at the end of the Visean, but in the Tarim Basin, it is still observed in the Serpukhovian. A similar situation is known in South China, where *Koninckopora* grainstones are observed in the Serpukhovian, Zone 18, Luocheng Formation, near Liuzhou, Guangxi (Anonymous, 1987).

5.4. The *Nanopora-Calcifolium* flora (Figs 5 A, D)

The *Calcifolium* algal microfacies is well documented in Skompski (1996). While *Nanopora* normally ranges from the Visean to the Serpukhovian (Fig. 5 A), *Calcifolium* (Fig. 5 D) is restricted to the latest Visean-Serpukhovian. Together, they are good markers for that level in the Tarim Basin, and they are quite common. Up to now, *Calcifolium* was known only from the occidental part of the Tethys, its easternmost report in Asia being the Tadzhikistan (Mamet, 1991). It has also been recognized in South China (Luocheng Formation, Qiaotoucun, Guangxi) (Mamet, unpublished).

5.5. The *Donezella* flora (Figs 6 B, C)

Donezella is a Paleosiphonocladale (Shuisky & Schirsova, 1985) that has caused considerable controversy (Riding 1979, Della Porta et al, 2002). As it is cosmopolitan and very abundant in the Northern Hemisphere, its role has intrigued many researchers and the taxon has been attributed to foraminifers, sponges, pseudosponges (Riding, 1979; Vachard, 1981). Although fragile, *Donezella* forms bafflestones, which usually crumple unless the thalli are

Figure 5. The following two figures illustrate widespread algae and characteristic algal microfacies. U. of M. stands for University of Montréal. A. *Nanopora fragilissima* (Maslov, 1939). Equatorial and longitudinal section in a well-sorted pellet-foraminifer (*Neoarchaediscus*) grainstone. U. of M. 924/12, Heshilafu Formation, Aitegou section, AET730, latest Visean Zone 16 (or slightly younger), X85. B. Diagenetically altered *Fasciella* strands on a brachiopod valve. U. of M. 924/10, as A, but X 22. C. *Paraepimastopora* sp. U. of M. 925/20, Tahaqi Formation, Paojianggou section, AET 355, Ghzelian, X 25. D. *Calcifolium okense* Shvetsov & Birina, 1935. U. of M. 924/11, as A. E. *Girvanella problematica* Nicholson & Etheridge, 1878. U. of M. 926/17, Atushi section BG 4, Visean, X85. F. *Petschoria elegans* Korde, 1951 (showing the lateral tubes). U. of M. 926/13, Lennan well 1, core 8, Bashkirian, X 54. G. *Koninckopora minuta* Weyer, 1968. U. of M. 924/21, Kelitake Formation, Aitegou section, AET 601, Early Visean Zone 11, X 25. H. *Rectangulina tortuosa* (Antropov, 1950). U. of M. 924/25, Kalawuyi Formation, Aitegou section, AET 745, Serpukhovian Zone 18, X70. I. *Claracrusta catenoides* (Homann, 1972). U. of M. 925/27, Qipan Formation, Qipan section, Kungurian (or younger), X 25. J. *Aciculella* sp. U. of M. 926/4, Manxi well 1, core 21, Asselian-Sakmarian, X 54. K. *Petschoria elegans* Korde, 1951 (showing the branching). U. of M. 926/10, Kalawuyi Formation, Lennan well 1, core 8, as F, but X 25. L. *Fourstonella? johnsoni* (Flügel, 1966). U. of M. 924/32, as H. M. *Beresella* ex. gr. *B. polyramosa* Kulik, 1964. U. of M. 925/9, Azigan Formation, Aitegou section, AET 774, Moscovian Zone 24, X25. N. *Pseudodonezella tenuissima* (Berchenko, 1982). U. of M. 924/27, as H. O. *Petschoria elegans* Korde, 1951 (showing the medullary tubes). U. of M. 926/12, Lennan well 1, core 8, Bashkirian, X 25. P. *Beresella erecta* Maslov & Kulik, 1956. U. of M. 925/8, Kalawuyi Formation, Aitegou section, AET 774, Moscovian Zone 24, X 35. Q. *Anthracoporellopsis machaevii* Maslov, 1956 (showing conceptacles). U. of M. 925/3, Kalawuyi Formation, Aitegou section, AET 754, Bashkirian? or early Moscovian? Zone 22-23, X54.

solidified by early marine cementation (see Figs 6 B, C). The sedimentary role of that algal incertae sedis has been recently discussed by Della Porta et al, 2002.

5.6. *The Petschoria flora (Figs 5 F, K, O)*

Petschoria has a wide dispersion. While it is known from the Urals, China, Yukon Territory and New Mexico, it is usually scarce. The taxon is well displayed in the Bashkirian-Moscovian limestone, well 1, Lennan (Figs 5 F, K, O).

5.7. *The Beresella-Dvinella flora (Figs 5 M, P)*

This duo is one of the most prolific “reef” builders (bafflestone) in Carboniferous-Early Permian time and is cosmopolitan in the Northern Hemisphere flora (Morin et al, 1994, Granier & Grgasovic, 2000). As the thalli have an outer mucilaginous coating fossilized as a continuous early cement layer, they are less susceptible than *Donezella* to crumple, and the bafflestones in the mud-mounds are usually better preserved. For taxonomy, refer to Granier & Grgasovic, 2000 or Forke & Samankassou, 2000 for upper Carboniferous microfacies. In the Tarim Basin, these bafflestones are restricted to the late Moscovian Azigan Formation.

5.8. *The Tubiphytes flora? (Fig. 6, F,G)*

This cosmopolitan algal? incertae sedis has been attributed to various phylla ranging from the hydrozoans to the cyanobacteria. Recent studies of Senowbarry-Daryan & Flügel (1993) suggest a polygenetic origin for this problematicum. It was illustrated from Permian Chinese carbonate sections by Mu (1982, 1985).

Whatever its origin (Vachard et al., 2001), *Tubiphytes* is a common encruster associated in the Tarim Basin with

bryozoans and sponges. It extends higher up associated with the *Ellesmerella* and *Pseudovermiporella* floras. In other parts of the world, *Tubiphytes* is usually associated with phylloid algae (*Eugonophyllum*, *Ivanovia*), but these are uncommon in the Tarim carbonates.

5.9. *The Ellesmerella flora (Fig. 6,E,H)*

Originally described from the Early Permian of the Canadian Arctic (Mamet et al., 1987), the flora has a great dispersion (Carnic Alps, Oman, Iran, China, Bolivia). It is probably the most prolific alga observed in the Permian limestones of the Tarim Basin. It is a common encruster of *Gymnocodium* (Fig.6,H).

5.10. *The Pseudovermiporella flora*

This is another controversial cosmopolitan alga. It has been excluded from the *Dasycladales* by Granier & Deloffre (1994) and transferred to the *Thaumatoporelles*. It has been repeatedly illustrated from China (Zhang & Wang, 1974; Zhao et al., 1981; Mu, 1981; Yang & Jiang, 1981; Mu, 1982; Yang et al., 1984; Lin, 1986; Wu, 1991, Riding & Guo, 1991; Sha et al., 1992).

6. Conclusions

The Late Paleozoic algal flora of the Tarim Basin is quite prolific. The most important taxa are “cosmopolitan” in the Northern Hemisphere. The only exception is that of *Calcifolium* which is reported here for the first time in the oriental part of the Tethys. Ten characteristic local algal assemblages are recognized from the Tournaisian to the Kungurian. They permit the establishment of a crude, local biostratigraphy at the stage level.





Figure 6. A. Well sorted pellet-calcisphere packstone with patches of grainstone. Calcispheres are mostly *Calcisphaera laevis* Williamson, 1881 with a few *Archaesphaera* sp. Foraminifers are conspicuously absent. Environment is a restricted, quiet lagoon. U. of M. 928/0, Kelitake Formation, Lennan well 50, core 9, Tournaisian, X 22. B. *Donezella* bafflestone. Intertwined, anastomosed network of *in situ* thalli, preserved by early marine isopachous cementation. Algae represent 15-20% of the volume and originally left an extensive porosity. Figured elements are scarce pellets, ostracods and molluscs. Marine facies, in euphotic zone, below wave action. U. of M. 926/8, Kalawuyi Formation, Tahe well 1, core 8, Bashikirian, X 22. C. Reworked *Donezella* bafflestone. Although there is little evidence of transport, the bafflestone (see B) is disarticulated and the thalli are fragmented. No evidence of early marine cementation. Marine facies, in euphotic zone, base of wave action zone. U. of M. 925/4, Kalawuyi Formation, Aitegou 1 section, AET 771, Moscovian, X 22. D. Strands of *Claracrusta catenoides* (Homann, 1972) encrusting fenestellid bryozoans fronds. Marine oncolite facies in high energy *Eoparafusulina* grainstone. U. of M. 925/33, Keziliqiman Formation, Duwa section, DW 34, Artinskian X22. E. *Ellesmerella* encrustations on dissolved mollusc shell. Marine oncolite facies in high-energy *Pseudoschwagerina* grainstone. U. of M. 925/32, Tahaqi Formation, Duwa section, DW 28, Asselian-Sakmarian, X22. F-G. Complex composite bindstones of *Tubiphytes obscurus* Maslov, 1956, fenestellid fronds, calcareous sponges, *Claracrusta catenoides* (Homann, 1972) strands and serpulopsids. *Tubiphytes* tissue encloses thin-shelled ostracods and sponge spicules. U. of M. 925/29 and 925/30, Qipan Formation, Qipan section, no number, Kungurian or younger, X 22. H. *Ellesmerella* encrustation on *Gymnocodium* in a *Sphaeroschwagerina* high-energy grainstone. *Ellesmerella* is the most abundant alga in the Early Permian of South China, while the *Gymnocodium* flora is usually developed in the upper part of the Permian. U. of M. 925/22, Tahaqi Formation, Paojianggou section, AET 375, Asselian-Sakmarian, X 22.

Acknowledgements. The authors would thank Prof. Yang Wanrong for providing some samples and Prof. Liao Weihua for helping to collect some references. This cooperative study is financially supported by Academia Sinica and by the Natural Sciences and Engineering Research Council of Canada.

7. References

- Anonymous, 1987. Carboniferous Carbonate Sequences in Guangxi. *11th International Congress of Carboniferous Stratigraphy and Geology*, Beijing. Guidebook, Excursion 6, 28p.
- ANTROPOV, I.A., 1950. New species of Devonian foraminifers of certain parts of the Russian Platform (in Russian, translated) *Akademiya Nauk SSSR, Kazanskoe Filial, Izvestia Seriya Geologicheskaya*, 1, 21-33.
- BERCHENSKO, O.I., 1982. *New calcareous algae species of the upper Serpukhovian deposits of the Donbass depression* (in Russian, translated) *Akademiya Nauk Ukrainkogo SSR, Institut geologicheskii Nauk, Naukova Dumka*, 51-55.
- DELLA PORTA, G., KENTER, F.A., & BAHAMONDE, F.R., 2002. Microfacies and paleoenvironment of *Donezella* accumulations across an Upper Carboniferous high-rising carbonate-platform (Asturias). *Facies*, 46, 149-168.
- FLÜGEL, E., 1966. Algen aus dem Perm der Karnischen Alpen. *Carinthia II*, 25, 76 pp.
- FORKE, H.C., & SAMANKASSOU, E., 2000. Biostratigraphic correlations of Late Carboniferous (Kasimovian) sections in the Carnic Alps : integrated paleontological data, facies and discussion. *Facies*, 42, 177-210.
- GRANIER, B. & DELOFFRE, R. 1994. Inventaire critique des algues Dasycladales fossiles. III partie. Les algues Dasycladales du Permien et du Trias. *Revue de Paléobiologie*, 14, 1, 49-84.
- GRANIER, B. & GRGASOVIC, T., 2000. Les Algues Dasycladales du Permien et du Trias. *Geologica Croatica*, 53, 1, 1-197.
- GROVES, J. R. & BRECKLE, P. L., 1997. Graphic correlation of Upper Paleozoic outcrop sections in the Western Tarim Basin, China. In Ross, C. A., Ross, J. R. & Breckle, P. L. (eds.) *Late Paleozoic Foraminifera: their biostratigraphy, evolution and paleoecology*. Cushman Foundation for foraminiferal Research, Special Publication, 36, 55-58.
- HOMANN, W., 1972. Unter- und tief-mittelpermische Kalkalgen aus den Rattendorfer Schichten, dem Trogkofel-Kalk der Karnischen Alpen (Oesterreich). *Senckenbergiana Lethaea*, 53, 135-313.
- Institute of Geology, Xinjiang Geological Bureau & Institute of Geology, CAGS, 1987. *The Carboniferous and Permian Stratigraphy and biota in Kalpin Region, Xinjiang*. Ocean Press, Beijing, 277p.
- JIA, Chengzhao, YIAO, Huijun, GAO, Jie, ZHOU, Dongyan & WEI, Guoqi, 1992. The Stratigraphy of Tarim Basin, In: Tong, Xiaoguang & Liang, Digang (eds.) *Symposium of Oil and Gas Exploration in Tarim Basin*. Xinjiang Technical and Health Publishing House, p.34-45.
- JIA, Chengzhao, WEI, Guoqi & LI, Liangsheng, 1992. The plate tectonic evolution and characters of main tectonic units in Tarim Basin, In: Tong, Xiaoguang & Liang, Digang (eds.) *Symposium of Oil and Gas Exploration in Tarim Basin*. Xinjiang Technical and Health Publishing House. p. 207-225.
- KORDÉ, K.B., 1951. New species and genera of calcareous algae from the Carboniferous of northern Urals (in Russian, translated). *Bulletin de la Société des Naturalistes de Moscou*, 1, 175-182.
- KULIK, E.L., 1964. Beresellids from the Carboniferous of the Russian Platform (in Russian, translated). *Paleontologicheskii Zhurnal*, 2, 99-114.
- LIAO, Wei Hua & RODRIGUEZ, S., 1999. Lower Carboniferous corals from the southwestern margin of the Tarim Basin, N.W. China, *Geobios*, 32, 4, 539-559.



- LIAO, Zhuting, WANG, Yujing, WANG, Keliang, JI-ANG, Nayan, XIA, Fengsheng, ZHOU, Yuxing, SUN, Fusheng & LI, Shoujun, 1992. Carboniferous of Tarim and its adjacents, In: Zhou Zhiyi and Chen Piji (eds.) *Biostratigraphy and Geological Evolution of Tarim*, Science Press, Beijing, p.202-242.
- LIN, Jiaying, 1986. Ecologic environment of the late Permian foraminifera fauna from the Liushe of Yishan, Guangxi, Zhuang Autonomous Region. Yichang Institute of Geology and Mineralogical Resources, *CAGS, Bulletin* 12, p. 91-99.
- MAMET, B., 1974. Une zonation par foraminifères du Carbonifère inférieur de la Téthys Occidentale. *Septième Congrès International de Stratigraphie et de Géologie du Carbonifère, Krefeld 1971, Compte Rendu*, 3, 391-408.
- MAMET, B., 1991. Carboniferous calcareous algae. In: R. Riding (ed.) *Calcareous algae and Stromatolites*, Springer-Verlag p.370-451.
- MAMET, B., 1998. A late Devonian microfossil with dasyclad algae affinities from northwestern Australia, *Alcheringa*, 22, 21-28.
- MAMET, B., ROUX, A. & NASSICHUK, W.W. 1987. Algues Carbonifères et Permiennees de l'Arctique Canadien. *Geological Survey of Canada, Bulletin* 342, p. 1-143.
- MASLOV, V.P., 1939. A discovery of Paleozoic green algae in Kazakhstan (in Russian, translated) *Problems of Paleontology*, 5, 290-296.
- MASLOV, V.P., 1956. *Fossil calcareous algae of the USSR* (in Russian, translated). Akademiya Nauk SSSR. Trudy Geologicheskogo Instituta, 160, 301pp.
- MASLOV, V.P. & KULIK, E.L., 1956. A new algal tribe (Bereselleae) in the carboniferous of the USSR (in Russian, translated). *Doklady Akademiya Nauk SSSR*, 106, 5, 126-129.
- MORIN F., DESROCHERS, A. & BEAUCHAMP, B., 1994. Facies analysis of Lower Permian platform carbonates, Sverdrup Basin, Canadian Arctic Archipelago, *Facies*, 31, 105-130.
- MU, Xinan, 1981. Upper Permian calcareous algae from western Guizhou. *Acta Palaeontologica Sinica*, 20, 1, 34-49.
- MU, Xinan, 1982. *Some calcareous algae from Xizang*. The Series of the Scientific expedition to the Qinghai-Xizang Plateau. Book V. Science Press, Beijing, p.205-240.
- MU, Xinan, 1985. *Calcareous algae of Late Carboniferous*, In: Mountaineering and Scientific Exploration Team, Academia Sinica ed. *Geology and Palaeontology In Mt. Tuomuur of Tianshan*. Xinjiang People's Publishing House.
- NAKAI, H. & KATO, M. 1981. On then occurrence of Koninckopora (Calcareous Alga) from the Viséan of Shikoku, Japan. *Journal Faculty of Science, Hokkaido University*, 19, 4, 446-447.
- NICHOLSON, H.A. & ETHERIDGE, R. Jr., 1878. A monograph of the Silurian fossils of the Girvan district, Ayrshire, with special reference to those contained in the "Gray Collection". Blackwood & Sons. Edinburgh and London, 341 pp.
- PINARD, S. & MAMET, B., 1998. Taxonomie des petits foraminifères du Carbonifère supérieur-Permien inférieur du bassin de Sverdrup, Arctique Canadien. *Palaeontographica Canadiana*, 15, p. 1-253.
- RIDING, R., 1979. *Donezella* bioherms in the Carboniferous of the southern Cantabrian Mountains, Spain. *Bulletin du Centre de Recherche Exploration-Production, Elf-Aquitaine*, 3, 2, 787-794.
- RIDING, R., & GUO, L., 1991. Permian marine calcareous algae. In: R. Riding (ed.), *Calcareous algae and Stromatolites*, Springer-Verlag, p. 452-480.
- SENOWBARI-DARYAN, B. & FLÜGEL, E., 1993. *Tubiphytes* Maslov, an enigmatic fossil: classification, fossil record and significance through time, part I: discussion of late Paleozoic material, In: Barattolo F. et al. (eds.) *Studies on Fossil Benthic Algae*, Mucchi, p. 353-382.
- SHA, Jingeng, ZHANG, Linxin, LUO, Hui, XU, Shan-hong & BAO, Huiming, 1992. On the closure age of the late Palaeozoic rift in Hohxii, Qinghai. *Acta Micropalaeontologica Sinica*, 9, 2, 177-182.
- SHUISKY, V. P. & SCHIRSCHOVA D. I. 1985. On the position of the Palaeoberesellids and other segmented algae from the Siphonophyceae. In: *New Data on the Geology, Biostratigraphy and Palaeontology of the Urals*. Akademia Nauk SSR, Uralskii Nauchnii Centr, Institut Geologii i Geochemii, p. 86-95.
- SKOMPSKI, ST., 1996. Stratigraphic position and facies significance of the limestone bands in the subsurface Carboniferous succession, Lublin Upland. *Acta Geologica Polonica*, 46, 3-4, 171-268.
- SHVETSOV, M.S., & BIRINA, L.M., 1935. On the petrology and origin of the Oka Limestone of the Mikhailov-Aleksin region (in Russian, translated) *Trudy Moskovskogo Geologicheskii Trust*, 10, 1-86.
- VACHARD, D., 1981. *Téthys et Gondwana au Paléozoïque Supérieur: les Données Afghanes*. Documents et Travaux de l'Institut Géologique Albert de Lapparent, 2, 463p.
- VACHARD, D., HAUSER, M., MARTINI, R., ZANINETTI, L., MATTER, A., & PETERS, T., 2001. New algae and problematica of algal affinity from the Permian of the Asselah Unit of the Batain Plain (East Oman). *Geobios*, 34, 4, 375-404.
- WANG, Hansheng, 1986. Tectonic evolution of Tarim Basin and direction of prospecting large scale oil/gas fields. *Xinjiang Geology*, 4, 3, 53-63.
- WANG, Qingming, NISHIDAI, T., & COWARD, M. P., 1992. The Tarim Basin NW China: Formation and aspects of petroleum geology. *Journal of Petroleum Geology*, 15, 5-34.
- WILLIAMSON, W.C., 1881. On the organization of the fossil plants of the Coal-Measures (Part X). *Philosophical Transactions of the Royal Society, London*, 171, 493-539.
- WRIGHT, P., 1981. Ultrastructure and early diagenesis of the Viséan alga *Koninckopora*. *Palaeontology*, 24, 1, 185-194.
- WU, Yasheng, 1991. Calcareous algae from Permian reef of Longlin, Guangxi, China. *Acta Palaeontologica Sinica*, 30, 6, 750-767.
- XIA, Guoying, DING, Yunjie & ZHAO, Songyin, 1987. The Subdivision of the Yuemengou group in Northern

- China, In: *Selected Papers on Carboniferous and Permian Coal-bearing Strata*, Geological Symposium, China. Science Press, Beijing, p. 9-17.
- YANG, Wanrong & JIANG, Nayan, 1981. On the depositional characters and microfacies of the Changhsing Formation and Permian-Triassic boundary in Changxing, Zhejiang. *Nanjing Institute of Geology and Palaeontology, Academia Sinica, Bulletin 2*, 113-130.
- YANG, Wanrong, ZANG, Quinlan & WANG, Ziyu, 1984. Petrological characteristics and environmental analysis of the biolithe from Chihhsia Formation of Mt. Xixia, Nanjing. *Journal of Stratigraphy*, 8, 1, 38-48.
- ZHANG, Linxin & WANG, Yujin, 1974. Permian algae, In: *Handbook of Stratigraphy and Palaeontology of Southwest China*. Nanjing Institute of Geology and Palaeontology, Academia Sinica, p. 315-317.
- ZHAI, G., & SUN, J., 1994. Structural evolution and distribution of oil and gas in the Tarim Basin. First International Symposium on the Petroleum Geology of the Tarim Basin, NW China, Stanford University, School of Earth Sciences, Stanford, California, Abstracts (unpaginated).
- ZHANG, Shibei & GU, Weiguo, 1991. Carboniferous and Lower Permian, In Zhang, Shibei & Gao, Qinqin (eds.) *Sinian to Permian Stratigraphy and Palaeontology of the Tarim Basin, Xingjiang: Part 2, Kalping-Bachu region*. The Petroleum Industry Press Beijing, p. 79-115.
- ZHAO, Jinke, SHENG, Jinzhang, YAO, Zhaoqi, LIANG, Xiluo, CHEN, Chuzhen, RUI, Lin & LIAO, Zhuoting 1981. The Changhsingian and Permian-Triassic boundary of South China. Nanjing Institute of Geology and Palaeontology, *Academia Sinica, Bull 2*, 1-112.
- ZHAO, Zhixin, HAN, Jianxiu & WANG, Zengji, 1984. *The Carboniferous strata and its fauna from Southwestern margin of Tarim Basin in Xinjiang*. Geological Publishing House Beijing, 187p.
- ZHU, Zili, 1995. The Carboniferous and Permian depositional sequences and fusulinid zonation in the Southwestern margin of the Tarim Basin, China. *XIII International Congress on Carboniferous-Permian Stratigraphy and Geology*, Krakow, Abstracts, p. 157.

Manuscript received 4.06.2003 ; accepted for publication 5.01.2004.