

CLIMATOLOGICAL CHARACTERISTICS OF NDVI TIME SERIES: CHALLENGES AND CONSTRAINTS

Stéphanie Horion^{1*}, Bernard Tychon² and Yves Cornet³

(1) Institute for Environment and Sustainability, Joint Research Centre of the European Commission, Ispra, Italy

(2) Département des Sciences et Gestion de l'Environnement, Université de Liège, Belgique

(3) Unité de Géomatique, Département de Géographie, Université de Liège, Belgique

* Corresponding author. E-mail address: stephanie.horion@jrc.ec.europa.eu

Abstract

Many studies already investigated the impact of climate change and climate variability on vegetation at global and continental scales. Low resolution satellite imagery is one of the main sources of information. In this paper, we describe a strategy to improve the quality of 10-daily time series of Normalized Difference Vegetation Index derived from SPOT-VEGETATION. A specific methodology was also defined in order to identify optimal test sites for the analysis of climate control on intra-annual dynamic of croplands. Finally two cases studies are presented to illustrate this research and in particular the non linear relationship between NDVI and meteorological parameters during the growing season.

Keywords

Vegetation dynamic, climate variability, low resolution imagery, geostatistics, time series analysis

Résumé

L'impact du changement climatique sur la végétation a déjà fait l'objet de nombreuses études à l'échelle continentale. L'imagerie satellitaire basse résolution est une importante source d'information pour ce genre d'étude. Cet article présente une méthodologie mise en place afin de maximiser l'information extraite de séries temporelles de l'indice normalisé de végétation (synthèses de 10 jours) pour l'analyse de l'impact de la variabilité climatique sur les cultures. La sélection des sites d'étude a également fait l'objet d'une grande attention afin de minimiser les facteurs perturbateurs du signal satellitaire autres que le climat. Pour terminer, deux cas d'étude seront présentés : le cas de la région d'Amhara en Ethiopie et celui de la Picardie en France.

Mots-clés

Dynamique de la végétation, variabilité climatique, image satellitaire basse résolution, géostatistique, analyse de séries temporelles

1. INTRODUCTION

Many studies already investigated the impact of climate change and climate variability on vegetation at global and continental scales. Using time series of remote sensing and climate data, Nemani *et al.* (2003) analysed trends in Net Primary Production in relation with changes in climate and showed that, between 1982 and 1999, primary productivity increased by 6% globally in response to climate change. This study also stressed the need to take into account the spatial variability of climatic constraints on plant growth when analysing the climate change impact on vegetation. Others authors described different phenomena linked with climate change, such as increases of seasonal Normalized Difference Vegetation Index amplitude and of growing season duration in the Northern high latitudes (Myneni *et al.*, 1998) or changes in circumpolar photosynthetic activities (Bunn *et al.*, 2005).

Understanding the interactions between climate and

vegetation is also a key issue in the PhD research of the main author (S. Horion) of this article. However, unlike other studies, we do not consider the vegetation globally but we focus on two specific types of vegetation: croplands and grasslands. One of our main objectives is to identify the meteorological factors that limit the development of croplands and grasslands in relation with their geographical localisation.

Because they provide both spatial and temporal data in large amount, low resolution satellites are customarily used as primary source of information on vegetation status and on meteorological conditions. However their coarse spatial resolution is greatly limiting their potential use. Indeed many vegetation types, land covers or natural processes can co-exist in a 1 km² pixel.

In this paper, we describe our strategy to improve the quality of 10-daily time series of Normalized Difference Vegetation Index derived from SPOT-VEGETATION. A specific methodology was also defined in order to identify optimal test sites for our research. Finally two




Figure 1. (left) Administrative layer: countries and regions, (right) Global Land Cover 2000, with legend simplified to 6 classes.

cases studies are presented to illustrate the non linear relationship between NDVI and meteorological parameters during the growing season.

2. DATA

2.1. Remote sensing Normalized Difference Vegetation Index

The Normalized Difference Vegetation Index (NDVI) is widely used as a proxy of the vegetation status and health. It has proven its efficiency in many research fields such as the monitoring of land cover changes and land degradation (Lambin *et al.*, 2001 ; Lambin, 2000), the crop growth monitoring and yield forecasting (Balaghi *et al.*, 2008 ; Zhang *et al.*, 2005), the study of vegetation phenology and its evolution over time (White *et al.*, 2006 ; Verstraete *et al.*, 2007), the interaction between climate and vegetation (Yu *et al.*, 2003 ; Vogt *et al.*, 2000 ; White *et al.*, 1997) and more specifically the impact of climate change on vegetation (Nemani *et al.*, 2003 ; Zhou *et al.* 2001 ; White *et al.*, 2005 ; Bunn *et al.*, 2005 ; Myneni *et al.*, 1998).

The NDVI is derived from the red and the near infrared bands following the equation:

$$NDVI = \frac{(NIR - RED)}{(RED + NIR)} \quad (\text{eq. 1})$$

Where NIR and RED are the amount of near infrared and red light reflected by the spectral object and recorded by the sensor (here: VEGETATION 1 and 2 sensors). This index ranges from -1 to +1 and relies on the absorption in the red part of the light spectrum due to the chlorophyll contained in the leaves, and on the scattering of the NIR by the mesophyll cells of the leaves. The more green and turgescient are the leaves, the closer to 1 is the NDVI. A null value represents more or less the threshold between the presence and the absence of vegetation.

For our research, ten years of 10-daily SPOT-VEGETATION NDVI were acquired from April 1998 till May 2008 for the entire globe. The original image data set has a spatial resolution of 1 km². However, before using

it for time series analysis, it went through a 3-steps procedure during which pixel values were aggregated to administrative entities (fig. 1a).

The first step is the pre-processing of raw images, i.e. geometric and atmospheric corrections and the computation of ten-days synthesis images, carried out by the Image Processing and Archiving Centre (CTIV, Mol) (Passot, 2000).

The next step is the computation of the NDVI regional unmixed statistics. Within a given region (fig. 1a), pixels covered totally by cropland are used for the calculation of the regional NDVI mean and standard deviation (Eerens *et al.*, 2004; Genovese *et al.*, 2001) (fig. 2). As reference for the land cover types, we used the Global Land Cover 2000 (Bartholomé *et al.*, 2005; Mayaux *et al.*, 2006) generalized in 6 classes at the original 1 km² resolution, GLC2000-6C (fig. 1b). This aggregation step is really important as it allows us to focus on a certain type of vegetation with a specific phenology.

Finally the last processing step is the temporal smoothing of the time series. During this step residual noise due to mainly undetected cloud/snow is filtered out.




Figure 2. Regional unmixed mean. In dark green, pixels considered to calculate the regional statistics.

2.2. Global meteorological data set

The global meteorological dataset used in this study has been acquired freely through the FOODSEC portal of the Joint Research Centre (<http://cidportal.jrc.ec.europa.eu/home/idp/thematic-portals/foodsec-imageserver/>). Out of the twelve indicators available, six

were downloaded for this study (table 1). A seventh indicator was computed using the precipitation and the potential evapotranspiration. It corresponds to a proxy of the climatic water balance. Time series are composed by 10-daily global images at 0.5 degree resolution. They are derived from 2 different sources, (1) the ERA40 re-analyses (Uppala *et al.*, 2005) for the period starting in January 1990 till August 2002 and (2) the operational

ECMWF (European Centre for Medium-Range Weather Forecast) atmospheric model for the period from September 2002 till May 2008. Regional statistics of the 7 meteorological indicators were also extracted per administrative region following a similar procedure than for the NDVI images, i.e. they were calculated using strictly the values of grid cells covered totally by agricultural areas in the GLC2000-6C (fig. 1b).

Table 1. 10-daily meteorological indicators.

Indicator	Unit	Abbreviation
Minimum temperature	°C	Tmin
Maximum temperature	°C	Tmax
Mean temperature	°C	Tmean
Cumulated precipitation	mm	Rain
Cumulated potential evapotranspiration	mm	ET0
Cumulated global radiation	KJ/m ² *day	Rg
Climatic water balance	mm	P-ETP

3. SITE SELECTION

Optimal test sites for the modelling of vegetation-climate interactions using the NDVI should ideally correspond to regions where the major part of the signal variation can be attributed to the climate and not to other phenomena such as land cover changes. For this reason our set of study cases has been established using a group of criteria which describe the spatial and temporal heterogeneities of the land covers and of the signal retrieved by the sensor (Horion *et al.*, 2007). The criteria and their implementation strategies are defined hereafter.

3.1. Spatial heterogeneity of land covers

The spatial heterogeneity introduced by the occurrence of different land covers in a region is evaluated with 2 different indices, the Relative Area and the Fragmentation index, both extracted from the generalized GLC2000-6C. The relative area is calculated for each land cover class existing in the administrative regions and gives information on the relative importance of land covers within the region. This indicator is important to assess the reliability of the regional NDVI statistics. Indeed the unmixing process used for the extraction of these statistics already improves the quality of the dataset by considering only pixels completely covered by croplands. But the regional statistics might nevertheless not be reliable or representative of the regional vegetation state if they are derived from too few pixels. The *Fragmentation index* refers to the spatial pattern in the surroundings of a pixel (3x3 window) (Eq. 2). The overall fragmentation of a region is evaluated by

averaging the Fr values of each pixel contained in that region.

$$F_r = \frac{(c_r - 1)}{(n_r - 1)} \quad (\text{eq. 2})$$

Where c_r is the number of different classes observed in a 3x3 window and n_r is the number of cells in the kernel (always 9 cells in this case).

3.2. Temporal heterogeneity of land covers (or land cover changes)

MODIS Land Cover products, MODIS LC, have been acquired in order to evaluate the error introduced by the use of a static land-cover map such as the GLC2000. These products are computed annually at a 1-km² resolution (Friedl *et al.*, 2002). The temporal trajectory of each pixel evaluated with MODIS LC for the period 2001-2004 is analyzed in order to identify land cover changes. Aberrant behaviours are identified using simple logical rules and such pixels are excluded from the land cover change analysis. A temporal trajectory is considered as aberrant or incoherent if one of the following rules is not respected: (1) a maximum of two land cover changes is acceptable in the temporal trajectory of the pixel; (2) the 'Shrublands' class is the only accepted phase prior to 'Forest'; (3) pixels with a temporal trajectory oscillating between water and another class are removed.

3.3. Spatial heterogeneity of the signal recorded by the sensor

The Coefficient of Variation CV measures the relative dispersion of a variable (Eq. 3). It is used here to esti-

mate how much the spectral signature of croplands is varying inside a same region.

$$CV_r = \frac{1}{d} \sum_{i=1}^d \left(\frac{Std_r}{Mean_r} \right) \quad (\text{eq. 3})$$

Where Std_r and $Mean_r$ are the standard deviation and average values calculated using the NDVI values of pixels covered by cropland and pertaining to region r : d is the 10-daily period (decade). The final CV_r corresponds to the average of all 10-daily $CV_{r,d}$.

4. RESULTS AND DISCUSSION

4.1. Optimal test sites for croplands

The analysis of the indicators presented in the previous section allowed us to identify candidate regions for our research. Figure 3a presents the relative area (RA) of cropland for our area of interest which comprises Europe and the part of Africa situated in the Northern Hemisphere. In Europe, regions mostly dominated by croplands ($RA > 60\%$) are quite numerous and spread over the entire area, notably in northern France, in Germany, in Poland, in Ukraine and in Spain. By contrast, in Africa, they are more concentrated on the Sahelian belt.

Like the relative area, the annual Fragmentation index per region Fr is also a good indicator of the spatial heterogeneity caused by mixed land covers. Most regions in our area of interest present a very low level of fragmentation, with Fr equal or below 0.125, meaning that in the 3×3 kernel (9 km^2) only two different land covers are observed in average (fig. 3b).

The temporal trajectories of the MODIS LC images from 2001 to 2004 have been analyzed to identify incoherencies and to locate the pixels where land cover changes occurred from 2001 to 2004. The identification of stable regions is crucial as we are working on the response of vegetation to climatic events. Using regions with a high land cover change dynamic or incoherent changes would affect our analysis by introducing extra-variability in the vegetation response retrieved from the sensor. Croplands are probably the most stable land cover in Europe, followed by the forest (fig.3c). The agricultural regions of France (Nord Pas-de-Calais, Picardie, Ile de France, etc) have recorded few land cover changes as well as the Pô region, the Garonne plain and Ukraine. On the contrary, regions with prevailing cropland stability are almost absent in Africa. Indeed land cover changes are important especially in the cropland and shrubland areas. Very few African regions can be




Figure 3. (top left) Relative Area of cropland derived from GLC2000, (top right) Spatial fragmentation of land covers, (bottom left) Land cover changes between 2001 and 2004, (bottom right) Coefficient of variation of NDVI for cropland (in grey, no data).




Figure 4. (left) Region of Amhara, Ethiopia; (right) Region of Picardie, France.

selected as potential agricultural regions for our study. The coefficient of variation is globally low ($CV < 0.2$), except in some areas in the Middle East and in Egypt (fig.3d). This suggests that the signal recorded at the sensor for the pixels covered by croplands can be considered homogeneous at the region scale.

4.2. Two cases studies

After evaluation of the preceding results, 15 regions were selected as test cases for our research, 10 in Europe and 5 in Africa. In this section we present two test cases, one in Ethiopia (fig.4a) and one in France (fig.4b).

The first case study corresponds to the region of Amhara in the Northern highlands of Ethiopia. Its size is about 165,000 km², of which more than 70 % are covered by cropland. The Coefficient of Variation of NDVI for cropland and the land cover fragmentation are both quite small ($CV_r = 0.22$; $Fr = 0.01$). However the percentage of stable cropland is rather low (about 11 %) and incoherent changes recorded between 2001 and 2004 by the MODIS Land Covers are important (12 %). Figure 5 (left) shows the typical landscape of Amhara. The agricultural fields are quite small (between 50 and 100 m in length) and mainly located on the plateau areas.

These observations and the importance of the relief in this area can explain the relatively high score for incoherent changes.

The second case corresponds to the region of Picardie in northern France. Its size is about 20,000km², of which more than 85 % are covered by stable croplands. The medium field size, much bigger than in Amhara, is about 500x100 m (fig.5b). Incoherent changes are negligible (< 2 % of the total area) and CV_r is also really low (~0.11), which indicates that the NDVI values recorded for all VEGETATION pixels covered by cropland in the region are similar.

Figure 6 presents the 10-daily evolution of NDVI for the two case studies in parallel with the 10-daily evolution of a limiting meteorological factor: 10-daily cumulated rainfall in the case of Amhara and mean temperature in the case of Picardie. The seasonality linked to the growing cycle of plants is obvious in both cases. In Ethiopia, a double crop cycle can be observed: from April to May-June (*Bleg* season) and from July to October (*Meher* season). In Picardie, the NDVI green-up is generally observed in March and the end of the growing season in October. The identification of a double crop season is less clear even though a small increase of the NDVI is often observed in August-September.



Figure 5. Google Earth sample images taken in the Amhara region (left) and in Picardie (right), with an identical scale.




Figure 6. 10-daily evolution of NDVI for cropland compared (top) with evolution of 10-daily cumulated rainfall in Amhara (Ethiopia) and (bottom) with evolution of 10-daily mean temperature in Picardie (France).

A good correlation between the NDVI cycle and the limiting meteorological parameter is also recorded in both regions. In Amhara, the rainfall increase precedes that of NDVI while in Picardie, cycles of NDVI and mean temperature are shifted in the opposite way. Scatterplots between NDVI and the limiting meteorological parameter show also interesting results regarding the type of relation between both variables (fig. 7a and 7b). We see clearly that fitting a linear regression line on the cloud of points is not a good solution. This demonstrates that a meteorological factor does not have the same impact on the NDVI evolution throughout the year and even through the growing season. In order to find statistically significant linear relation which is also meaningful from an agro-meteorological point of view, we need to split the growing season into separate (phenological) phases. For example, in the case of Picardie, a good linear relation can be found between March and June ($R^2 = 0.65$), which corresponds respectively to the greening phase and to the beginning of the maturing process for winter crops (USGS crop calendar, [\[www.usda.gov/oce/weather/CropCalendars/index.htm\]\(http://www.usda.gov/oce/weather/CropCalendars/index.htm\), consulted in December 2009\).](http://</p>
</div>
<div data-bbox=)

Following those preliminary results, a detailed investigation of interrelations between meteorological events and the vegetation response is currently undergone with specific focuses on (1) identifying key phenological stages within the growing season, (2) considering the cumulated impact of meteorological events (e.g. rainfall deficit during several decades, heat wave, etc) and (3) considering the possibility of a delayed response of vegetation.

5. CONCLUSION AND PERSPECTIVES

We have presented here some results of the PhD research of S. Horion concerning the impact of climate on croplands using two low resolution global datasets, respectively 10-daily NDVI estimated for cropland and 10-daily meteorological parameters combining ERA-40 and ECMWF data (table 1).

A specific strategy was set up to improve these data-




Figure 7. Scatterplots between 10-daily NDVI and a limiting meteorological parameter: (left) 10-daily cumulated rainfall in Amhara (Ethiopia), (right) mean temperature in Picardie (France).

sets and to focus only on agricultural areas. The Global Land Cover 2000 was used to identify VEGETATION pixels completely covered by cropland. These pixels were then used to compute regional statistics of NDVI and meteorological parameters.

The next step was the selection of optimal test cases for the analysis of climate control on croplands. Ideally, NDVI seasonal signal of such regions should be mainly influenced by the variation of meteorological parameters. Therefore regions with high fragmentation in landscapes, high land cover change dynamics and high spatial variation of NDVI with respect to the mean signal were excluded. The final choice was based on the percentage of stable cropland cover.

Further work will be dedicated to the analysis of cross-correlations between the monthly NDVI and the 7 meteorological parameters. As the two case studies showed, interactions between meteorological parameters and NDVI can not be modelled using a simple linear regression over the year or the growing season. Indeed they need to be studied at a time scale smaller than the growing season in order to identify properly the limiting factors on plant growth. Like the limiting factors, which are variable from a region to another, the time scale for the analysis needs also to be adjusted for each region, taking into account the phenology of the vegetation under consideration. Moreover, we also considered in our analysis the possibility of a delayed response of the vegetation and/or a cumulated effect of meteorological events (up to 3 months).

Acknowledgments

The PhD research of S. Horion was founded by the F.R.I.A (Fonds pour la formation à la Recherche dans l'Industrie et dans l'Agriculture). We would like also to thank the Agriculture Unit (European Commission, Joint Research Centre) and Herman Eerens (VITO) for

providing us with regional unmixed statistics of SPOT-VEGETATION NDVI.

References

- Balaghi, R., Tychon, B., Eerens, H. and Jlibene, M., 2008. Empirical regression models using NDVI, rainfall and temperature data for the early prediction of wheat grain yields in Morocco. *International Journal of Applied Earth Observation and Geoinformation*, 10(4), 438-452.
- Bartholomé, E. and Belward, A., 2005. GLC2000: a new approach to global land cover mapping from Earth observation data. *International Journal of Remote Sensing*, 26(9), 1959-1977.
- Bunn, A. G., Goetz, S. J. and Fiske, G. J., 2005. Observed and predicted responses of plant growth to climate across Canada. *Geophysical Research Letters*, 32, L16710.
- Eerens, H., Piccard, I., Royer, A. and Orlandi, S., 2004. Methodology of the MARS Crop Yield Forecasting System (Vol.3). Remote sensing information, data processing and analysis. Eds. Royer, A. and Genovese, G. European Commission, DGJRC. EU 21291/EN3. 53-56.
- Friedl, M.A., McIver, D.K., Hodges, J.C.F., Zhang, X.Y., Muchoney, D. Strahler, A.H., Woodcock, C.E., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F. and Schaaf, C., 2002. Global land cover mapping from MODIS: algorithms and early results. *Remote Sensing of Environment*, 83(1-2), 287-302.
- Genovese, G., Vignolles, C., Nègre, T. and Passera, G., 2001. A methodology for a combined use of normalised difference vegetation index and CORINE land cover data for crop yield monitoring and forecasting. A case study on Spain. *Agronomie*, 21, 91-111.
- Horion, S., Eerens, H., Tychon, B. and Cornet, Y., 2007. Spatial aggregation of low resolution satellite data for the monitoring of vegetation response to climatic stresses: Analysis of the spatial heterogeneity of aggregated entities. European Space Agency, (Special Publication) ESA SP-636, July 2007, 6p.
- Lambin, E., 2000. Land Cover Assessment and Monitoring. in Wiley J.:Encyclopedia of Analytical Chemistry. Eds. R.A. Meyers, John Wiley and Sons Ltd, Chichester, 22p.
- Lambin, E.F., Turner, II B.L., Geist, H., Agbola, S., Angelsen, A., Bruce, J.W., Coomes, O., Dirzo, R., Fischer, G., Folke, C., George, P.S., Homewood, K., Imbernon, J., Leemans, R., Li, X., Moran, E.F.,

- Mortimore, M., Ramakrishnan, P.S., Richards, J.F., Skånes, H., Steffen, W., Stone, G.D., Svedin, U., Veldkamp, T., Vogel, C. and Xu, J., 2001. The Causes of Land-Use and Land-Cover Change: Moving beyond the Myths. *Global Environmental Change*, 11, 261-269.
- Mayaux, P., Strahler, A., Eva, H., Herold, M., Shefali, A., Naumov, S., Dorado, A., Di Bella, C., Johansson, D., Ordoyne, C., Kopin, I., Boschetti, L. and Belward, A., 2006. Validation of the Global Land Cover 2000 Map. *IEEE Trans. on Geoscience and Remote Sensing*, 44(7), 1728-1739.
- Myneni, R.B., Tucker, C.J., Asrar, G. and Keeling, C.D., 1998. Interannual variations in satellite-sensed vegetation index data from 1981 to 1991. *Journal of Geophysical Research*, 103, 6145-6160.
- Nemani, R.R., Keeling, C.D., Hashimoto, H., Jolly, W.M., Piper, S.C., Tucker, C.J., Myneni, R.B. and Running, S.W., 2003. Climate-driven increases in global terrestrial net primary production from 1982 to 1999. *Science*, 300, 1560-1563.
- Passot, X., 2000. VEGETATION image processing methods in the CTIV. In *VEGETATION 2000: 2 years of operation to prepare the future*. Ed. Saint G., Lake Maggiore, Italy, 15-21.
- Uppala, S.M., Kallberg, P.W., Simmons, A.J., Andrae, U., Bechtold, V.D., Fiorino, M., Gibson, J.K., Haseler, J., Hernandez, A., Kelly, G.A., Li, X., Onogi, K., Saarinen, S., Sokka, N., Allan, R.P., Andersson, E., Arpe, K., Balmaseda, M.A., Beljaars, A.C.M., Van De Berg, L., Bidlot, J., Bormann, N., Caires, S., Chevallier, F., Dethof, A., Dragosavac, M., Fisher, M., Fuentes, M., Hagemann, S., Holm, E., Hoskins, B.J., Isaksen, L., Janssen, P.A.E.M., Jenne, R., McNally, A.P., Mahfouf, J.F., Morcrette, J.J., Rayner, N.A., Saunders, R.W., Simon, P., Sterl, A., Trenberth, K.E., Untch, A., Vasiljevic, D., Viterbo, P. and Woollen, J., 2005. The ERA-40 re-analysis. *Quarterly Journal of the Royal Meteorological Society*, 131, 2961-3012.
- Verstraete, M.M., Gobron, N., Aussedat, O., Robustelli, M., Pinty, B., Widlowski, J.-L. and Taberner, M., 2007. An automatic procedure to identify key vegetation phenology events using the JRC-FAPAR products. *Advances in Space Research*, 41, 1773-1783.
- Vogt, J.V., Niemeier, S., Beaudin, I. and Viau, A., 2000. Drought Monitoring from Space. *Advances in Natural and Technological Hazards Research*, 14, 167-183.
- White, M.A. and Nemani, R.R., 2006. Real-time monitoring and short-term forecasting of land surface phenology. *Remote Sensing of Environment*, 104(1), 43-49.
- White, M.A., Hoffman, F., Hargrove, W.W. and Nemani, R.R., 2005. A global framework for monitoring phenological responses to climate change. *Geophysical Research Letters*, 32, L04705.
- White, M.A., Thornton, P.E. and Running, S.W., 1997. A continental phenology model for monitoring vegetation responses to interannual climatic variability. *Global Biogeochemical Cycles*, 11, 217-234.
- Yu, F., Price, K.P., Ellis, J. and Shi, P., 2003. Response of seasonal vegetation development to climatic variations in eastern central Asia. *Remote Sensing of Environment*, 87, 42-54.
- Zhang, P., Anderson, B., Tan, B., Huang, D. and Myneni, R. B., 2005. Potential monitoring of crop production using a satellite-based Climate-Variability Impact Index. *Agricultural and Forest Meteorology*, 132, 344-358.
- Zhou, L., Tucker, C.J., Kaufmann, R.K., Slayback, D., Shabanov, N.V. and Myneni, R.B., 2001. Variations in northern vegetation activity inferred from satellite data of vegetation index during 1981 to 1999. *Journal of Geophysical Research*, 106(D17), 20069-20083.
-